

cell-matrix interactions as determined by transendothelial electrical resistance measurement. Therefore, we hypothesized that 4-HNE-induced permeability changes involved modulation of focal adhesion and adherens junction proteins. Treatment of BLMVECs with 4-HNE resulted in the redistribution of FAK,  $\beta$ -catenin, paxillin, VE-cadherin, and ZO-1 and caused intercellular gap formation. Western blot analyses confirmed that 4-HNE formed Michael adducts with the focal adhesion and adherens junction proteins. Furthermore, 4-HNE decreased tyrosine phosphorylation of FAK without affecting total cellular FAK contents. Flow cytometry and fluorescent microscopy analyses revealed a time-dependent reduction in the surface integrins after 4-HNE treatment. **Conclusion:** These results indicate that 4-HNE affects EC permeability by modulating cell-cell adhesion involving focal adhesion, adherens and tight junction proteins, as well as integrins signal transduction.

Supported by NIH RO1 HL 69909 and P01 HL 58064 to V.N.

## 46

**HYPERCAPNIA INDUCES MUCIN GENE EXPRESSION IN HUMAN AIRWAY EPITHELIAL CELLS.** N. Wang, M.C. Niend, A. Nair, J.J. Sznajder, P.H.S. Sporn, Pulmonary and Critical Care Medicine, Feinberg School of Medicine, Northwestern University, Chicago, IL.

**Objective:** Mucus hypersecretion is a prominent feature of chronic obstructive pulmonary disease, asthma, and cystic fibrosis. These diseases may be associated with development of hypercapnia, acutely during exacerbations and/or chronically in their advanced stages. In this study, we tested the hypothesis that hypercapnia might contribute to mucus overproduction by increasing expression of genes for mucins produced by airway epithelial cells. **Methods:** The human mucocutaneous carcinoma cell line (NCI-H292) was used in this study. Cells were cultured in the presence of 5% CO<sub>2</sub> (control; equivalent to PCO<sub>2</sub> ≈ 40 mm Hg) or 20% CO<sub>2</sub> (hypercapnia; equivalent to PCO<sub>2</sub> ≈ 160 mm Hg), in the absence or presence of phorbol 12-myristate 13-acetate (PMA; 10 nM), an inflammatory agent known to induce mucin gene expression, for 6 to 96 hours. mRNA for the mucin genes *MUC2*, *MUC5AC*, *MUC5B*, and *MUC19* were quantitated by real-time PCR and normalized to expression of 18S ribosomal RNA. **Results:** In the absence of PMA, culture of NCI-H292 cells in 20% CO<sub>2</sub>, as compared to 5% CO<sub>2</sub>, increased expression of *MUC5AC* by ≈ 3-fold. In comparison, PMA (in 5% CO<sub>2</sub>) increased *MUC5AC* mRNA expression ≈ 70-fold. Culture of NCI-H292 cells in 20% CO<sub>2</sub> augmented the PMA-induced increase in *MUC5AC* mRNA levels by ≈ 3-fold as compared to stimulation with PMA in 5% CO<sub>2</sub>. The increases in *MUC5AC* mRNA in response to 20% CO<sub>2</sub> peaked at 24 hours. *MUC19* mRNA expression was increased in the presence of 20% CO<sub>2</sub> in the absence and presence of PMA to a degree similar to that for *MUC5AC*. Neither 20% CO<sub>2</sub> nor PMA increased expression of mRNA for *MUC2* or *MUC5B*. **Conclusion:** Hypercapnia stimulates mucin gene expression in vitro and may be important in driving gene expression and synthesis of mucins in vivo. These results reveal a previously unrecognized stimulus of mucin gene expression. Furthermore, they suggest that correcting hypercapnia may be an important strategy for reducing mucus hypersecretion in patients with acute and chronic lung diseases.

## 47

**INCREASED NOX1 IN NEOINTIMAL SMOOTH MUSCLE CELLS IS ASSOCIATED WITH REDOX-SENSITIVE SIGNALING AND ACTIVATION OF MATRIX METALLOPROTEINASE 9.** S. Xu,<sup>1</sup> D. Jagadeesha,<sup>2</sup> A. Chamseddine,<sup>1</sup> R. Bhalla,<sup>2</sup> F. Miller Jr,<sup>1</sup> <sup>1</sup>Internal Medicine, <sup>2</sup>Anatomy and Cell Biology, University of Iowa, Iowa City, IA.

The vascular response to injury involves redox-dependent activation of smooth muscle cells (SMC) and development of a neointima. Activation of matrix metalloproteinase 9 (MMP-9) appears to be critical in the development of arterial lesions. We hypothesized that increased NADPH oxidase-derived superoxide (O<sub>2</sub><sup>-</sup>) by neointimal SMC contributes to activation of MMP-9. Two weeks after balloon injury of rat aorta, neointimal SMC showed higher O<sub>2</sub><sup>-</sup> levels compared with adjacent medial SMC. A role for NADPH oxidase was suggested by the observations that neointimal SMC O<sub>2</sub><sup>-</sup> levels were reduced by the flavoenzyme inhibitor DPI, and immunostaining for p47<sup>phox</sup> and p22<sup>phox</sup> were increased in the neointima, as compared with medial, SMC. Neointimal and medial SMC were isolated and grown in culture. Consistent with the findings in situ, O<sub>2</sub><sup>-</sup> levels and NADPH oxidase activity were significantly greater in neointimal SMC. Nox1 mRNA levels were more than 2-fold increased in neointimal SMC, whereas Nox4 expression was similar to that of medial SMC. ERK1/2 kinase activation was greater in neointimal vs medial SMC and was inhibited by tiron and DPI. MMP-9 activity, as measured by gelatin zymography, was higher in neointimal SMC compared to medial SMC in basal conditions and 48 hours after IL-1 $\beta$  stimulation and was reduced by pretreatment with the ERK inhibitor PD98059 or DPI. We conclude that NADPH oxidase-derived O<sub>2</sub><sup>-</sup> signals activation of MMP-9 following vascular injury. Moreover, Nox1-dependent cell signaling in neointimal SMC may play an important role in the pathogenesis of vascular disease.

Supported by AHA and NIH 14388 and HL081750.

## 48

**THE SILENCING MEDIATOR OF RETINOID AND THYROID HORMONE RECEPTORS REPRESSES ADIPOGENESIS VIA ITS DISTAL INTERACTING DOMAIN.** C. Yu, M.M.

Sutanto, R.N. Cohen, Department of Medicine, The University of Chicago, Chicago, IL. Obesity is known to be a risk factor for the development of type 2 diabetes mellitus and the metabolic syndrome. Thus, it is important to understand the factors that modulate adipogenesis. Adipogenesis requires the nuclear hormone receptor (NHR) peroxisome proliferator-activated receptor gamma (PPARGamma). NHRs recruit coactivators (CoAs) in the presence of agonists to increase gene transcription; in addition, some NHRs also recruit nuclear receptor corepressors (CoRs) under certain circumstances. CoRs, such as the nuclear corepressor protein (NCoR) and the silencing mediator of retinoid and thyroid hormone receptors (SMRT), recruit a histone deacetylase complex to repress gene transcription. Using a RNA interference system to decrease SMRT and NCoR expression in 3T3-L1 cells, we have previously shown that CoRs repress adipogenesis and decrease the expression of adipocyte-specific proteins by modulating PPARGamma transcriptional activity. We have additionally shown that the intracellular balance of CoRs and CoAs dictates the ability of thiazolidinediones (TZDs), which are PPARGamma ligands used clinically in the treatment of diabetes mellitus, to stimulate adipocyte differentiation. To define the mechanisms of CoR recruitment by PPARGamma in 3T3-L1 cells, we have now focused on the corepressor SMRT. SMRT contains two interacting domains (IDs) that are required for interactions with NHRs. Interestingly, and contrary to other NHRs, PPARGamma interacts with SMRT specifically via the SMRT C-terminal ID (SMRT ID1). To define the amino acids

important in specifying the ability of ID1 to function, residues within this domain were mutated to the corresponding residues in the proximal SMRT ID (ID2). Interestingly, a single amino acid within ID1, which is not conserved in ID2, appears to be vital in the recruitment of SMRT by PPARGamma. In contrast, this specific sequence is not required for SMRT interactions with the retinoic acid receptor (RAR), another NHR that interacts strongly with SMRT. Thus, the ability of PPARGamma to recruit the corepressor SMRT is dependent on specific residues within the SMRT C-terminal ID that are distinct from amino acids important for interactions with other NHRs. We have shown that the ability of TZDs to up-regulate PPARGamma-mediated processes is dependent on the balance of CoRs and CoAs. These data also suggest that it might be possible to specifically alter PPARGamma-SMRT interactions to modify TZD action on adipogenesis.

## 49

**NOVEL ROLE OF PRE-B-CELL COLONY-ENHANCING FACTOR IN PULMONARY ENDOTHELIAL BARRIER REGULATION.** L.Q. Zhang, J. Cepeda, J.G.N. Garcia, S.Q. Ye, Section of Pulmonary/Critical Care, Department of Medicine, The University of Chicago, Chicago, IL.

**Rationale:** Our prior genomic and genetic studies identified pre-B-cell colony-enhancing factor (PBEF) as a potential novel biomarker in acute lung injury (ALI) (Ye et al, 2005). To elucidate the molecular mechanism underlying PBEF in the pathogenesis of ALI, we assessed the role of PBEF in in vitro vascular barrier regulation using confluent human pulmonary artery endothelial cell (HPAEC) or microvascular cells (HMVEC-L) monolayers since increased vascular permeability is a cardinal feature of ALI. **Methods:** Inhibition of PBEF expression was achieved by the PBEF siRNAs. Overexpression of the PBEF gene was accomplished by adeno-PBEF Endothelial cells (EC) were transfected with PBEF siRNA or infected with adeno-PBEF for 48 hours before treated with thrombin (0.1 U/mL) or S1P (1 nM) for various time. Parameters to evaluate endothelial cell barrier functions include transendothelial electric resistance (TER), actin staining, myosin light chain phosphorylation, calcium influx. **Results:** Reductions in PBEF protein expression (> 70%) by siRNA significantly attenuated EC barrier dysfunction induced by the potent edemagenic agent, thrombin reflected by reductions in TER (60 to 70% reduction). Furthermore, PBEF siRNA blunted thrombin-mediated increases in increases in Ca<sup>2+</sup> entry, polymerized actin formation, and myosin light chain phosphorylation, events critical to the thrombin-mediated permeability response. Overexpression of PBEF by adeno-PBEF vector significantly increases EC permeability reflected by decreased TER in HMVEC-L by ≈ 30%, indicating that overexpression of PBEF results in dysfunction. **Conclusions:** These in vitro observations strongly support that PBEF is critically involved in the endothelial cell barrier regulation. It encourages us to further explore the mechanistic insight into PBEF in the murine model of ALI.

Funded by NHLBI SCCOR-Molecular Approaches to Ventilator-Associated Lung Injury (U01HL 073994).

## 50

**SPHINGOSINE 1-PHOSPHATE DRAMATICALLY ALTERS THE HUMAN PULMONARY ARTERY ENDOTHELIAL CELLS' LIPID RAFT PROTEOME.** J. Zhao, P. Singleton, S. Dudek, J.G.N. Garcia, Department of Medicine, The University of Chicago, Chicago, IL.

**Rationale:** Lipid rafts are plasma membrane microdomains involved in dynamic membrane signaling and trafficking. Our previous studies (Singleton et al, FASEB J, 2005) have demonstrated that the platelet-derived lipid sphingosine 1-phosphate (S1P) plays a critical regulatory role in maintenance and enhancement of pulmonary vascular barrier function. S1P ligates the S1P receptor 1, which we have previously localized to lipid rafts of human pulmonary artery endothelial cells (HPAECs). Here we use 2-D electrophoresis analysis to characterize changes in tyrosine phosphoprotein content in lipid rafts of HPAECs challenged with S1P. **Methods and Results:** HPAECs were challenged with 1  $\mu$ M of S1P for 5 minutes, washed with cold PBS twice, and solubilized with 1% Triton X-100 at 4°C. The Triton X-100 insoluble material was subjected to discontinuous Optiprep™ gradient centrifugation and the light density (lipid raft) fractions were collected. Isolated lipid rafts were resolubilized in 50  $\mu$ L of 7 M urea/2 M thiourea and 50 mM DTT. Isoelectric focusing was carried out on a nonlinear electrophoresis gel strip (pH3–10, 7 cm) with the second dimension carried out on 4–20% SDS-PAGE gel. Analytical gels were stained using Imperial blue (Pierce) and Western blotting was performed using antiphosphotyrosine antibody. Protein levels were not significantly altered in lipid rafts isolated from control vs S1P-treated cells (1  $\mu$ M, 5 minutes). The 2-D gel images revealed over 30 proteins recruited to lipid rafts after S1P treatment (5 minutes) with antiphosphotyrosine blots showing S1P to increase tyrosine phosphorylation of over 20 proteins, including the src kinase family and cortactin. **Conclusions:** These results suggest that tyrosine kinases participate in S1P-induced signaling in lipid rafts and that 2-D proteomic analysis is a powerful tool for studying regulation of pulmonary endothelial cells barrier function.

Supported by NIH grant HL58064.

## 51

**HIGH-MOLECULAR-WEIGHT POLYETHYLENE GLYCOL PROTECTS THROMBIN-INDUCED ENDOTHELIAL BARRIER DYSFUNCTION BY INDUCING ACTIN CYTOSKELETON REARRANGEMENT, WHICH RESULTS IN ROBUST ENHANCEMENT OF ENDOTHELIAL CELL BARRIER INTEGRITY.** E.T. Chiang, S.M. Camp, S.M. Dudek, O. Zaborina, J.C. Alverdy, J.G. Garcia, The University of Chicago, Chicago, IL.

ARDS is characterized by sudden, life-threatening lung failure with diffuse alveolar infiltrate, reduced arterial oxygenation, and pulmonary edema. Acute lung injury (ALI) is the end result of common pathways initiated by a variety of local or systemic insults. Polyethylene glycol (PEG) compounds are inert and nontoxic polymers that act as a surrogate mucin lining providing protection against bacterial infections on intestinal epithelial cells. We hypothesized that PEG may provide similar protective effects on lung endothelium by attenuating endothelial cell (EC) activation that results in barrier dysfunction. This study examines the effects of a high-molecular-weight PEG (PEG15–20) on cultured human pulmonary microvessel EC from barrier disruptive agents. EC were cultured on microelectrodes and changes in transendothelial electrical resistances (TER) were measured to assess alterations in paracellular permeability. PEG induced a rapid, dose-dependent increase in TER similar to barrier-enhancing lipid such as sphingosine 1-phosphate (S1P). Optimal concentration of 7.5–9% PEG induced a robust increase in TER from 2,500 to 5,000 ohm, which was sustained for 40 hours. In comparison to S1P, which induced a 40% increase in resistance in 10–15 minutes, the maximum barrier enhancing effect of PEG was achieved