



OPEN ACCESS

Methylation in the matrix metalloproteinase-2 gene is associated with cerebral ischemic stroke

Hsiu-Fen Lin,^{1,2} Edward Hsi,^{3,4} Ling-Chun Huang,¹ Yi-Chu Liao,^{5,6}
Suh-Hang H Juo,^{3,4,7,8} Ruey-Tay Lin^{1,2}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jim-2016-000277>).

For numbered affiliations see end of article.

Correspondence to

Dr Ruey-Tay Lin, Department of Neurology, Kaohsiung Medical University, 100 TzYou First Road, Kaohsiung City 807, Taiwan; tay@kmu.edu.tw and Dr. Suh-Hang Hank Juo, Institute of New Drug Development, China Medical University, 91 Hsueh-Shih Road, Taichung City 404, Taiwan; hjuo@mail.cmu.edu.tw

Accepted 25 January 2017
Published Online First
13 February 2017

Copyright © 2016 American Federation for Medical Research

ABSTRACT

Matrix metalloproteinase-2 (MMP-2) is involved in the pathophysiology of stroke. Previous studies have shown that MMP-2 activity is increased in stroke; however, evidence of epigenetic regulation of the MMP-2 in stroke is still limited. We examined methylation of the MMP-2 promoter in patients with ischemic stroke. This study included 298 patients with ischemic stroke and 258 age-matched and sex-matched controls. MMP-2 promoter methylation levels were measured by pyrosequencing at eight potential cytosine-guanine (CpG) sites. Multivariate regression analysis was used to adjust for general stroke risk factors, and the specific effects of sex and stroke subtype were analysed. The methylation levels of MMP-2 in the peripheral blood of the patients with stroke were lower than controls in all eight CpG sites, especially at site 1, site 5, site 7, and site 8 (adjusted $p=0.036$, 0.002 , 0.021 , and 0.041 , respectively). In subgroup analysis by sex, a significant association was found only in men but not in women. When the stroke subtype was considered, men with small-vessel stroke had significantly lower methylation levels at all MMP-2 CpG sites than the controls (3.01% vs 3.65%, adjusted $p=0.018$). Although men with large-artery atherosclerosis stroke also had lower MMP-2 methylation levels, no significant difference was found (3.25% vs 3.65%, adjusted $p=0.253$). Demethylation of the MMP-2 promoter in patients with ischemic stroke was in a sex and stroke subtype-specific manners. These findings may add to the understanding of epigenetic modification of MMP-2 on ischemic stroke.

INTRODUCTION

Stroke is a leading cause of death and disability worldwide, and matrix metalloproteinases (MMPs), which are involved in extracellular matrix remodeling, have been reported to play a key role.^{1,2} The MMPs are a family of over 20 proteinases, in which gelatinase A (matrix metalloproteinase-2 (MMP-2)) is one of the main constitutive enzymes in the brain. MMP-2 has attracted increasing interest in the field of stroke pathophysiology,^{3,4} and it has been demonstrated that MMP-2 is involved in the atherosclerotic process.⁵ Higher levels of circulating MMP-2 have also been reported in cerebral ischemia,⁶ intracranial hemorrhage,⁷ and cerebral white matter hyperintensity.⁸ In

Significance of this study

What is already known about this subject?

- DNA methylation plays a major role in ischemic stroke.
- Matrix metalloproteinase-2 (MMP-2) activity is increased in cerebral stroke.
- The MMP-2 gene expression is regulated by epigenetic mechanisms, including methylation.

What are the new findings?

- This is the first study demonstrated lower MMP-2 methylation level in patients with ischemic stroke than controls.
- This effect was especially pronounced in men.
- With regards to the stroke subtype, MMP-2 methylation appeared to have more obvious influence in small-vessel stroke than large-vessel stroke.

How might these results change the focus of research or clinical practice?

- These findings may add to the understanding of epigenetic modification of MMP-2 on ischemic stroke.

addition, previous studies have suggested that MMP-2 plays multiple roles in ischemic stroke, including participating in the injury process in the early stage and contributing to recovery during the later stage.^{9,10} In the human brain, levels of MMP-2 may remain high in ischemic lesions for up to several years after stroke.¹¹ Taken together, these findings suggest that MMP-2 is a crucial regulatory marker for cerebral ischemic stroke.

Emerging evidence suggests that DNA methylation plays a major role in ischemic stroke.^{12,13} In adult cells, DNA methylation takes place in a cytosine-guanine (CpG) dinucleotide context, also known as a CpG site, and in general, DNA methylation silences gene expression.¹⁴ It is known that the expression of the MMP-2 gene is regulated by epigenetic mechanisms,^{5,15} and it has also been demonstrated that a decrease in methylation of the MMP-2 gene can result in an increase in its



CrossMark

To cite: Lin H-F, Hsi E, Huang L-C, et al. *J Investig Med* 2017;**65**:794–799.

expression in diseases including lung cancer, pterygium, and atherosclerosis.^{5 16 17} Although previous studies have shown that circulating MMP-2 levels are increased in cerebral ischemia,⁶ no published study has investigated the relationship between MMP-2 methylation and ischemic stroke.

In this study, we evaluated methylation levels in the MMP-2 promoter using DNA from the peripheral blood of patients with ischemic stroke. Sex has been reported to be an important effect modifier regarding stroke risk factors and outcomes,¹⁸ and our previous study showed that global methylation had different effects on men and women with stroke.¹⁹ Therefore, we also investigated possible sex-specific effects on MMP-2 methylation in patients with ischemic stroke. As an increase in MMP-2 activity after ischemic stroke has only been reported in the small-vessel subtype of stroke,⁶ we also investigated MMP-2 methylation levels in patients with different stroke subtypes.

MATERIALS AND METHODS

Subjects

Stroke subjects

Two hundred and ninety-eight patients with ischemic stroke, aged between 40 and 80 years were selected from our previously reported patient cohort,²⁰ all of whom were recruited from Kaohsiung Medical University Hospital and Taichung Veterans General Hospital in Taiwan. These patients were chosen as follows: (1) those with stroke in whom the first stroke episode occurred before 80 years of age; (2) these patient were then split into four age groups (40–50 years, 51–60 years, 61–70 years, and 71–80 years); and (3) an equal number of men and women were randomly selected from each age group. All of these patients were diagnosed with ischemic stroke according to the WHO criteria,²¹ which included laboratory examinations and cranial CT or MRI. Ischemic stroke was diagnosed when brain imaging showed acute infarction or revealed no evidence of hemorrhage. Stroke subtypes were classified according to the Trial of ORG 10172 in Acute Stroke Treatment (TOAST).²²

Control subjects

The healthy controls were recruited from those who answered an advertisement at Kaohsiung Medical University Hospital and who reported no history of stroke or myocardial infarction.²³ Two hundred and fifty-eight age-matched and sex-matched controls were recruited using random number tables created by SPSS statistical software (SPSS, Chicago, Illinois, USA). To minimise effects of

age and gender, we selected an equal number of male and female cases and controls from each age group.

Data on socio-demographics, medical history of hyperlipidemia, hypertension, diabetes, and cigarette smoking were obtained for each participant. All the study subjects (including cases and controls) had no cancer history or peripheral basement membrane remodeling disease by self-report. Glucose, triglycerides, and total cholesterol levels were tested from venous blood after fasting for at least 8 hours. Subjects with stroke had their blood collected in a subacute or chronic stroke stage. None of the samples were collected within 72 hours after acute stroke. A current smoker was defined as anyone who had smoked within the past 12 months of enrollment, and an ex-smoker was defined as someone who had stopped smoking for more than 1 year. Hypertension was diagnosed as systolic or diastolic blood pressure $\geq 140/90$ mm Hg or receiving antihypertensive medication. Diabetes was diagnosed as fasting blood glucose ≥ 126 mg/dL or receiving treatment for diabetes. Hypercholesterolemia was diagnosed as a serum level of total cholesterol ≥ 200 mg/dL or receiving lipid-lowering medications. The Institutional Review Boards of Kaohsiung Medical University Hospital and Taichung Veterans General Hospital approved this study, and the participants provided written informed consent.

DNA extraction and MMP-2 methylation level measurement

Genomic DNA was extracted using a commercially available DNA extraction kit (Gentra; Qiagen, Hilden, Germany). DNA samples were then treated with sodium bisulfate to convert unmethylated cytosine (C) to uracil and leave methylated C intact using an EpiTect Fast Bisulfite Kit (Qiagen) following the manufacturer's recommendations.

Using PyroMark Assay Design V2.0 software (Qiagen), the MMP-2 methylation assay was designed to cover eight CpG sites in the promoter of the MMP-2 gene. The bisulfite-modified DNA was used to amplify the 178-bp product in the MMP-2 promoter area (figure 1) (primers are shown in online supplementary table). Quantitative MMP-2 methylation levels at the CpG sites were then measured by pyrosequencing (PyroMark Q24; Qiagen). Universal unmethylated and methylated DNA was run as the controls. Using PyroMark Q24 2.0.6 software (Qiagen), methylation quality was checked and quantified. The level of MMP-2 methylation was expressed as a percentage of methylated cytosine in methylated and unmethylated cytosines.




Figure 1 Schematic diagram for the design of the matrix metalloproteinase-2 methylation (MMP-2) assay. *F, R, and S denote forward, reverse, and sequence primers, respectively. CpG, cytosine-guanine sites.

Statistical analysis

The χ^2 test and Student's t-test were applied to analyse the characteristics of the study subjects. Associations between MMP-2 methylation level and ischemic stroke were analysed using the Student's t-test. Multivariate logistic regression analysis was then used to adjust for age, sex, smoking, hypertension, hypercholesterolemia, and diabetes. Data were also analysed by sex to assess sex-specific effects. The different effects of stroke subtypes were also assessed. SPSS software V18.0 for Windows was used for all statistical analyses (SPSS), and a two-tailed $p < 0.05$ was considered to be statistically significant.

RESULTS

The demographic characteristics of the study subjects are presented in [table 1](#). With regards to risk factors for stroke, 76.2% of the patients with stroke had hypertension, 51.3% had diabetes, 50.3% had hypercholesterolemia, and 35.6% were current or ex-smokers. In the control group, 45.7% had hypertension, 10.9% had diabetic, 31.4% had hypercholesterolemia, and 15.1% were current or ever smokers. All risk factors were significantly different between the cases and controls as expected. With regards to stroke TOAST classification, 18.8% of the cases had large-artery atherosclerosis, 11.7% had cardioembolism, 45.0% had small-vessel occlusion, and 24.5% had an undetermined etiology.

The methylation levels in the peripheral blood at the eight potential MMP-2 promoter CpG sites are shown in [table 2](#). In the MMP-2 CpG sites overall, the methylation levels were significantly lower among the patients with ischemic stroke compared with the controls in univariate analysis. After adjusting for traditional risk factors, the p values for CpG site 1, site 5, site 7, and site 8 were still significant (adjusted $p = 0.036$, 0.002 , 0.021 , and 0.041 , respectively). The average methylation level of all CpG sites was also significantly lower in the patients with ischemic stroke than in the controls (adjusted $p = 0.036$).

To assess sex-specific effects, we further stratified the participants by gender. Compared with the male controls, the male patients with ischemic stroke had lower MMP-2

Table 1 Demographic characteristics of the study participants

	Stroke n=298	Control n=258	p Value
Age (years)	61.5±10.3	61.6±10.5	0.901
Male (%)	147 (49.3)	121 (46.9)	0.610
Hypertension (%)	227 (76.2)	118 (45.7)	<0.001
Diabetes (%)	153 (51.3)	28 (10.9)	<0.001
Hypercholesterolemia (%)	150 (50.3)	81 (31.4)	<0.001
Current and ex-smoker (%)	106 (35.6)	39 (15.1)	<0.001
Stroke subtype			
Large-artery atherosclerotic	56 (18.8)		
Cardioembolism	35 (11.7)		
Small vessel	134 (45.0)		
Undetermined	73 (24.5)		

Data are shown as mean±SD for quantitative variables and n (%) for qualitative variables.

Table 2 Association between ischemic stroke and matrix metalloproteinase-2 methylation level

CpG position/ methylation %	Stroke n=298 Mean±SD	Control n=258 Mean±SD	Crude p value	Adjusted p value
Site 1	1.88±1.02	2.13±1.11	0.005	0.036
Site 2	3.75±1.97	4.24±2.29	0.006	0.052
Site 3	4.23±1.95	4.66±2.08	0.013	0.093
Site 4	3.59±1.84	3.96±2.10	0.028	0.103
Site 5	1.92±0.88	2.25±1.20	<0.001	0.002
Site 6	3.43±1.78	3.86±2.14	0.009	0.098
Site 7	2.70±1.17	3.02±1.41	0.003	0.021
Site 8	4.46±2.23	4.98±2.38	0.009	0.041
Average of all sites	3.24±1.52	3.64±1.74	0.004	0.036

Bold signifies $p < 0.05$.

Adjusted p values were adjusted for age, sex, hypertension, diabetes, hypercholesterolemia, and smoking status.

CpG, cytosine-guanine sites.

methylation levels at all CpG sites than the controls, and especially at CpG site 5 (adjusted $p = 0.011$) ([table 3](#)). While the methylation levels were also lower at all CpG sites in the female patients than in the female controls, none of the differences reached statistical significance.

Since previous studies have reported an association between MMP-2 plasma levels and the small-vessel stroke subtype rather than large-artery atherosclerosis,⁶ we further analysed MMP-2 methylation levels in men stratified by stroke subtype. Compared with the controls, the male patients with small-vessel stroke had a significantly lower average methylation level at all MMP-2 CpG sites than the controls (3.01% vs 3.65%, adjusted $p = 0.018$) ([figure 2](#)). Although the male patients with large-artery atherosclerosis also had a lower MMP-2 methylation level than the controls, the difference was not statistically significant (3.25% vs 3.65%, adjusted $p = 0.253$).

DISCUSSION


In this study, we found a lower MMP-2 methylation level in the DNA from peripheral blood of patients with ischemic stroke than in controls with no history of stroke. This effect was especially pronounced in men. With regards to the stroke subtype, MMP-2 methylation appeared to have a more pronounced effect in patients with small-vessel stroke than large-vessel stroke, which is consistent with a previous study that reported an increased MMP-2 serum level in patients with small-vessel stroke.⁶ To the best of our knowledge, this is the first study to investigate the association between methylation in the MMP-2 promoter and cerebral ischemic stroke, and in particular with regards to sex and stroke subtype-specific effects. The findings of this study add to the current knowledge of the epigenetic modification of MMP-2 on ischemic stroke.

MMP-2 has been reported to play a dual role in ischemic stroke.^{9 10} After a focal cerebral ischemic insult, the early pathological effect of MMP-2 is the disruption of the blood-brain barrier through degradation of the basal lamina, which may contribute to hemorrhagic transformation and eventually neuronal cell death.⁹ Conversely,

Table 3 Association between ischemic stroke and matrix metalloproteinase-2 methylation level by sex

CpG position/ methylation %	Men				Women			
	Stroke n=147 Mean±SD	Control n=121 Mean±SD	Crude p value	Adjusted p value	Stroke n=151 Mean±SD	Control n=137 Mean±SD	Crude p value	Adjusted p value
Site 1	1.81±0.98	2.14±1.13	0.012	0.064	1.94±1.06	2.12±1.10	0.153	0.253
Site 2	3.61±1.99	4.21±2.34	0.024	0.121	3.89±1.95	4.27±2.25	0.121	0.245
Site 3	4.08±1.97	4.73±2.18	0.011	0.135	4.38±1.94	4.59±2.00	0.354	0.453
Site 4	3.43±1.77	3.95±2.24	0.034	0.130	3.75±1.89	3.97±1.98	0.341	0.464
Site 5	1.82±0.83	2.25±1.31	0.001	0.011	2.02±0.91	2.25±1.10	0.051	0.061
Site 6	3.30±1.79	3.92±2.33	0.014	0.175	3.56±1.76	3.82±1.97	0.238	0.352
Site 7	2.62±1.15	3.04±1.48	0.010	0.059	2.77±1.19	3.01±1.35	0.109	0.210
Site 8	4.33±2.25	4.88±2.45	0.056	0.266	4.59±2.21	5.06±2.32	0.079	0.092
Average of all sites	3.12±1.51	3.65±1.87	0.011	0.095	3.36±1.53	3.64±1.61	0.141	0.221

Adjusted p values were adjusted for age, hypertension, diabetes, hypercholesterolemia, and smoking status.
CpG, cytosine-guanine sites.

**Figure 2** The association between promoter methylation level of matrix metalloproteinase-2 gene and stroke subtype in men.

MMP-2 also plays a role in endogenous repair, particularly in angiogenesis and in reestablishing cerebral blood flow during the poststroke recovery stage.¹⁰ Using gelatin zymography of human brain tissue, an increase in the activity of MMP-2 was observed in patients who died even several months after stroke.¹¹ This prolonged elevation implies that MMP-2 is involved in recovery after a stroke. Furthermore, several studies have investigated the correlation between MMP-2 genetic polymorphisms and ischemic stroke, and the results showed that the patients with risk alleles had a 1.5–2.6-fold higher risk of ischemic stroke.^{24–26} Taken together, these findings suggest that MMP-2 plays a crucial role in ischemic stroke.

Accumulating clinical and experimental evidence supports that DNA methylation is involved in the pathogenesis of ischemic stroke and that it is a risk factor for atherosclerosis, which is thought to be a major pathological change in stroke.^{27–28} In addition, several human studies have investigated associations between the level of DNA methylation and ischemic stroke.^{12–19} Moreover, in our previous study,¹⁹ we found lower global methylation levels in long interspersed nucleotide element-1 (LINE-1) in male patients with stroke than in controls. In our recent study,

we found that female patients with stroke had lower methylation levels of the estrogen receptor α gene than controls, especially in those with large-artery atherosclerosis and cardioembolic subtypes.¹² The genetic expression of MMP-2 is known to be epigenetically regulated.^{5–15} Although several studies have explored MMP-2 genetic methylation status in several diseases,^{5–16–17} no previous study has reported the effect of MMP-2 methylation in patients with ischemic stroke.

The clinical features of cerebral ischemic stroke including traditional risk factors and outcomes differ between men and women.^{18–29} Moreover, differences in plasma MMPs activity between men and women have been reported in human carotid plaque.³⁰ In our previous study, we found that menopausal women carrying an MMP-9 risk allele were predisposed to stiffer arteries, whereas this risk allele had no such effect on premenopausal women.³¹ In addition, a previous study reported that the expression of MMP-2 is closely related to sex hormones.³² These biological mechanisms and findings from other studies further support our finding that MMP-2 methylation has a sex-specific effect in patients with ischemic stroke.

It has been reported that the effects of MMP-2 are limited to specific stroke subtypes.^{6–24} Lucivero *et al*⁶ reported an increase in plasma MMP-2 level in patients with lacunar stroke, but not in those with large-artery atherosclerosis stroke. In addition, several MMP-2 polymorphisms were reported to be independent risk factors for small-vessel infarction in a Caucasian population, but not for other stroke subtypes.²⁴ A recent study on a Chinese population also reported a significant association between MMP-2 genetic polymorphism and small-vessel disease.²⁶ In the present study, a decrease in MMP-2 methylation level was only observed in the patients with small-vessel stroke, and not in those with large-artery atherosclerotic stroke, which is consistent with the previous results. Blood–brain barrier dysfunction has been reported to be a potentially important mechanism for small-vessel stroke,³³ and MMP-2 has been reported to play a role in disruption of the blood–brain barrier after cerebral hypoperfusion.³⁴ This could be a possible explanation for our finding of the association between MMP-2 demethylation and small-vessel stroke.

There are limitations to this study. It is likely that some healthy controls might still have other hidden vascular diseases. Therefore, our results may need to be cautiously interpreted. Another limitation is that we do not have record on stroke severity for further analysis. It is always a concern when using a blood marker for a cerebral disease. In the current study, we measured MMP-2 methylation levels in peripheral leucocyte DNA, however, we did not have data to support a consistent level of methylation between peripheral blood and the brain, nor samples to measure serum MMP-2 levels. Although there is lack of comprehensive data, our results are consistent with MMP-2 activity reported in previous clinical stroke studies.⁶ Further studies are warranted to further investigate MMP-2 methylation in the context of ischemic stroke.

In summary, this study demonstrates that male patients with ischemic stroke have a lower MMP-2 methylation level, and that this effect is especially pronounced in patients with small-vessel stroke. Future studies on developing strategies to alter the activity of MMP-2 in patients with ischemic stroke should take sex and stroke subtype-specific effects into consideration.

Author affiliations

¹Department of Neurology, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

²Department of Neurology, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

³Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

⁴Department of Medical Research, China Medical University Hospital, Taichung, Taiwan

⁵Department of Neurology, Taipei Veterans General Hospital, Taipei, Taiwan

⁶Department of Neurology, National Yang-Ming University School of Medicine, Taipei, Taiwan

⁷Graduate Institute of Biomedical Sciences, China Medical University, Taichung, Taiwan

⁸Institute of New Drug Development, China Medical University, Taichung, Taiwan

Contributors H-FL, S-HHJ, R-TL conceived and designed the experiments. Y-CL, EH performed the experiments. H-FL, EH, L-CH analysed the data. H-FL, S-HHJ, R-TL wrote the paper.

Funding This work was supported by grants from the Ministry of Science and Technology (Taiwan, R.O.C. 103-2314-B-037-029, 103-2314-B-037-027-MY2, and 104-2314-B-037-030), and Kaohsiung Medical University Hospital intramural grants (KMUH102-2R48).

Competing interests None declared.

Patient consent Obtained.

Ethics approval The Institutional Review Boards of Kaohsiung Medical University Hospital and Taichung Veterans General Hospital.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Romanic AM, White RF, Arleth AJ, *et al.* Matrix metalloproteinase expression increases after cerebral focal ischemia in rats: inhibition of matrix metalloproteinase-9 reduces infarct size. *Stroke* 1998;29:1020–30.
- Rosenberg GA, Navratil M, Barone F, *et al.* Proteolytic cascade enzymes increase in focal cerebral ischemia in rat. *J Cereb Blood Flow Metab* 1996;16:360–6.
- Cunningham LA, Wetzel M, Rosenberg GA. Multiple roles for MMPs and TIMPs in cerebral ischemia. *Glia* 2005;50:329–39.
- Yong VW, Power C, Forsyth P, *et al.* Metalloproteinases in biology and pathology of the nervous system. *Nat Rev Neurosci* 2001;2:502–11.
- Chen KC, Wang YS, Hu CY, *et al.* OxLDL up-regulates microRNA-29b, leading to epigenetic modifications of MMP-2/MMP-9 genes: a novel mechanism for cardiovascular diseases. *FASEB J* 2011;25:1718–28.
- Lucivero V, Prontera M, Mezzapesa DM, *et al.* Different roles of matrix metalloproteinases-2 and -9 after human ischaemic stroke. *Neurol Sci* 2007;28:165–70.
- Alvarez-Sabin J, Delgado P, Abilleira S, *et al.* Temporal profile of matrix metalloproteinases and their inhibitors after spontaneous intracerebral hemorrhage: relationship to clinical and radiological outcome. *Stroke* 2004;35:1316–22.
- Corbin ZA, Rost NS, Lorenzano S, *et al.* White matter hyperintensity volume correlates with matrix metalloproteinase-2 in acute ischemic stroke. *J Stroke Cerebrovasc Dis* 2014;23:1300–6.
- Heo JH, Lucero J, Abumiya T, *et al.* Matrix metalloproteinases increase very early during experimental focal cerebral ischemia. *J Cereb Blood Flow Metab* 1999;19:624–33.
- Silletti S, Kessler T, Goldberg J, *et al.* Disruption of matrix metalloproteinase 2 binding to integrin alpha v beta 3 by an organic molecule inhibits angiogenesis and tumor growth in vivo. *Proc Natl Acad Sci USA* 2001;98:119–24.
- Clark AW, Krekoski CA, Bou SS, *et al.* Increased gelatinase A (MMP-2) and gelatinase B (MMP-9) activities in human brain after focal ischemia. *Neurosci Lett* 1997;238:53–6.
- Lin HF, Hsi E, Liao YC, *et al.* Demethylation of circulating estrogen receptor alpha gene in cerebral ischemic stroke. *PLoS ONE* 2015;10:e0139608.
- Schweizer S, Meisel A, Marschenz S. Epigenetic mechanisms in cerebral ischemia. *J Cereb Blood Flow Metab* 2013;33:1335–46.
- Grumt I, Pikaard CS. Epigenetic silencing of RNA polymerase I transcription. *Nat Rev Mol Cell Biol* 2003;4:641–9.
- Pereira IT, Ramos EA, Costa ET, *et al.* Fibronectin affects transient MMP2 gene expression through DNA demethylation changes in non-invasive breast cancer cell lines. *PLoS ONE* 2014;9:e105806.
- Moran A, Fernandez-Marcelo T, Carro J, *et al.* Methylation profiling in non-small cell lung cancer: clinical implications. *Int J Oncol* 2012;40:739–46.
- Riau AK, Wong TT, Lan W, *et al.* Aberrant DNA methylation of matrix remodeling and cell adhesion related genes in pterygium. *PLoS ONE* 2011;6:e14687.
- Niewada M, Kobayashi A, Sandercock PA, *et al.* Influence of gender on baseline features and clinical outcomes among 17,370 patients with confirmed ischaemic stroke in the international stroke trial. *Neuroepidemiology* 2005;24:123–8.
- Lin RT, Hsi E, Lin HF, *et al.* LINE-1 methylation is associated with an increased risk of ischemic stroke in men. *Curr Neurovasc Res* 2014;11:4–9.
- Lin HF, Tsai PC, Liao YC, *et al.* Chromosome 9p21 genetic variants are associated with myocardial infarction but not with ischemic stroke in a Taiwanese population. *J Investig Med* 2011;59:926–30.
- The World Health Organization MONICA Project (monitoring trends and determinants in cardiovascular disease): a major international collaboration. WHO MONICA project principal investigators. *J Clin Epidemiol* 1988;41:105–14.
- Adams HP Jr, Bendixen BH, Kappelle LJ, *et al.* Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in acute stroke treatment. *Stroke* 1993;24:35–41.
- Lin HF, Liu CK, Liao YC, *et al.* The risk of the metabolic syndrome on carotid thickness and stiffness: sex and age specific effects. *Atherosclerosis* 2010;210:155–9.
- Fatar M, Stroick M, Steffens M, *et al.* Single-nucleotide polymorphisms of MMP-2 gene in stroke subtypes. *Cerebrovasc Dis* 2008;26:113–19.
- Nie SW, Wang XF, Tang ZC. Correlations between MMP-2/MMP-9 promoter polymorphisms and ischemic stroke. *Int J Clin Exp Med* 2014;7:400–4.
- Zhang M, Zhu W, Yun W, *et al.* Correlation of matrix metalloproteinase-2 single nucleotide polymorphisms with the risk of small vessel disease (SVD). *J Neurol Sci* 2015;356:61–4.
- Chan GC, Fish JE, Mawji IA, *et al.* Epigenetic basis for the transcriptional hyporesponsiveness of the human inducible nitric oxide synthase gene in vascular endothelial cells. *J Immunol* 2005;175:3846–61.
- Wang YS, Chou WW, Chen KC, *et al.* MicroRNA-152 mediates DNMT1-regulated DNA methylation in the estrogen receptor alpha gene. *PLoS ONE* 2012;7:e30635.

- 29 Sheikh K, Bullock CM. Effect of measurement on sex difference in stroke mortality. *Stroke* 2007;38:1085–7.
- 30 Hellings WE, Pasterkamp G, Verhoeven BA, *et al*. Gender-associated differences in plaque phenotype of patients undergoing carotid endarterectomy. *J Vasc Surg* 2007;45:289–96; discussion 296–287.
- 31 Lin RT, Chen CH, Tsai PC, *et al*. Sex-specific effect of matrix metalloproteinase-9 functional promoter polymorphism on carotid artery stiffness. *Atherosclerosis* 2012;223:416–20.
- 32 Natoli AK, Medley TL, Ahimastos AA, *et al*. Sex steroids modulate human aortic smooth muscle cell matrix protein deposition and matrix metalloproteinase expression. *Hypertension* 2005;46:1129–34.
- 33 Topakian R, Barrick TR, Howe FA, *et al*. Blood–brain barrier permeability is increased in normal-appearing white matter in patients with lacunar stroke and leucoaraiosis. *J Neurol Neurosurg Psychiatr* 2010;81:192–7.
- 34 Nakaji K, Ihara M, Takahashi C, *et al*. Matrix metalloproteinase-2 plays a critical role in the pathogenesis of white matter lesions after chronic cerebral hypoperfusion in rodents. *Stroke* 2006;37:2816–23.