

Paradigms in chronic obstructive pulmonary disease: phenotypes, immunobiology, and therapy with a focus on vascular disease

Michael Schivo,^{1,2} Timothy E Albertson,^{1,3} Angela Haczku,^{1,2}
 Nicholas J Kenyon,^{1,2} Amir A Zeki,^{1,2} Brooks T Kuhn,¹ Samuel Louie,^{1,2}
 Mark V Avdalovic^{1,3}

¹Department of Internal Medicine, University of California Davis School of Medicine, Sacramento, California, USA

²Center for Comparative Respiratory Biology and Medicine, Genome and Biomedical Sciences Facility, University of California Davis, Davis, California, USA

³Department of Medicine, Veterans Administration Northern California Healthcare System, Mather, California, USA

Correspondence to

Dr Michael Schivo,
 Department of Internal Medicine, University of California Davis School of Medicine, 4150 V Street, Suite 3400, Sacramento, CA 95817, USA;
 mschivo@ucdavis.edu

Accepted 9 February 2017
 Published Online First
 3 March 2017

Copyright © 2017 American Federation for Medical Research

ABSTRACT

Chronic obstructive pulmonary disease (COPD) is a complex and heterogeneous syndrome that represents a major global health burden. COPD phenotypes have recently emerged based on large cohort studies addressing the need to better characterize the syndrome. Though comprehensive phenotyping is still at an early stage, factors such as ethnicity and radiographic, serum, and exhaled breath biomarkers have shown promise. COPD is also an immunological disease where innate and adaptive immune responses to the environment and tobacco smoke are altered. The frequent overlap between COPD and other systemic diseases, such as cardiovascular disease, has influenced COPD therapy, and treatments for both conditions may lead to improved patient outcomes. Here, we discuss current paradigms that center on improving the definition of COPD, understanding the immunological overlap between COPD and vascular inflammation, and the treatment of COPD—with a focus on comorbid cardiovascular disease.

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) represents a major global health burden that is widely recognized as a complex, heterogeneous syndrome rather than a single disease. Epidemiological data reveal three major themes: first, COPD consists of several clinical phenotypes, most of which need further refinement in definition. By understanding these COPD phenotypes, it may be possible to improve treatment. Second, COPD is often recognized as a chronic inflammatory lung disorder with important immunological mechanisms and systemic manifestations. Appreciating the immunobiology of COPD may facilitate better treatment paradigms and shed light on common mechanisms shared between COPD and cardiovascular disease. Third, COPD often exists with and may potentiate cardiovascular disease independent of tobacco smoking. How COPD treatment affects cardiovascular disease, and vice versa, is also unclear.

In this review, we aim to: (1) discuss current COPD phenotypes based on relevant epidemiological biomarker studies; (2) review COPD

immunobiology with a focus on the overlap with cardiovascular disease; and (3) discuss recent advances in COPD treatment, including treatments that can affect both COPD and cardiovascular disease.

COPD PHENOTYPES

Decades ago, medical schools taught the concept that COPD existed as two basic clinical phenotypes: chronic bronchitis versus pulmonary emphysema. We currently understand that COPD is far more heterogeneous. The Global Initiative for Chronic Obstructive Lung Disease (GOLD) took steps to categorize COPD with greater sophistication beginning in 2001.¹ GOLD has subsequently taken the established staging of COPD by spirometry, primarily forced expiratory volume in 1 s as a percent of forced vital capacity (FEV₁/FVC or FEV₁%), and created patient groups that include evaluation of the burden of symptoms and exacerbation frequency in a more comprehensive assessment of the impact of COPD on patient lives. These groups include patients with good lung function and minimal symptoms (GOLD A) to patients with advanced lung disease and a high degree of symptoms (GOLD D). As expected, it also includes a group of patients with COPD with advanced decline in lung function but with relatively few symptoms (GOLD C), and a group with preserved lung function but a high degree of symptoms (GOLD B). Group B and C patients were not a surprise to clinicians, and having an empiric approach to categorizing these complex phenotypes of COPD was welcome.

Large prospective clinical cohort studies have improved our understanding of the heterogeneity of COPD. We review and highlight major discoveries that have emerged from these studies with particular emphasis on phenotyping schemes, contribution of CT scans, and the relationship of COPD with comorbid conditions, including cardiovascular disease.

Cohort studies

COPDGene

COPDGene was originally designed to be an observational study to identify genetic factors



CrossMark

To cite: Schivo M, Albertson TE, Haczku A, et al. *J Investig Med* 2017;**65**:953–963.

associated with COPD,² but it was refined to be a prospective cohort study enrolling 4500 smoker controls between 2008 and 2011 at 21 different clinical centers: 1500 GOLD stage 1, and 4500 GOLD stages 2–4 (total 10,500 subjects). Patients who were classified as ‘smoker controls’ had an FEV₁/FVC of >0.70 and a FEV₁ >80%, all post-bronchodilator. A small group of non-smoker controls were also included as a comparison for the quantitative CT scan data. Additionally, an interesting subcohort emerged labeled GOLD-U, which was an unclassified COPD cohort of smokers with a decrease in FEV₁ but a preserved FEV₁/FVC ratio.² The study goals were to characterize each of these groups with respect to symptoms, medications, and spirometry; inspiratory and expiratory CT scans; exercise capacity; and genome-wide association patterns to compare within each of these cohorts. At final enrollment, two-thirds of the subjects were non-Hispanic whites while one-third were African-American. The defining contribution of COPDGene is taking existing clinical staging (GOLD) and defining novel phenotypes within these stages.

ECLIPSE

The Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) was the first large prospective cohort designed to characterize COPD with the goal of discovering novel biomarkers.^{3,4} ECLIPSE enrolled patients over a 3-year period of time including 2164 patients with COPD and 582 control subjects (of whom 337 were smokers). Patients were assessed at eight different time points with the following studies: PFTs (including body plethysmography, spirometry and forced oscillometry, but not carbon monoxide diffusing capacity), biomarkers (including exhaled breath condensate (EBC)), clinical health outcomes (eg, death and disability), CT scans, body impedance, oxygen saturation, and 6 min walk distance. The advantage of the approach in ECLIPSE, as compared with the COPDGene, is that it took the basic definition of COPD and sought to define new phenotypes from that starting point over a 3-year period, for example, the ‘frequent exacerbator’.

MESA-LUNG

The Multi-Ethnic Study of Atherosclerosis (MESA) is a prospective cohort study that was designed to study the prevalence and progression of subclinical cardiovascular disease.⁵ The MESA cohort enrolled a total of 6814 subjects between the ages of 45 and 84 from six separate clinical sites across the USA. One of the recruitment emphases was to include a highly multiethnic cohort, and the participants included non-Hispanic whites, Hispanics, African-Americans, and Asians. MESA-LUNG is a nested study that uses the data from MESA to test a specific hypothesis: that endothelial dysfunction plays a specific role in the pathogenesis of COPD and, more specifically, emphysema.⁶ Initially, MESA-LUNG recruited 3965 randomly sampled participants from MESA that included 24% African-Americans, 23% Hispanics, and 18% Chinese-Americans. These patients had spirometry, quantitative CT scan data, as well as a wide range of genetic and biometric data. MESA-LUNG defined cardiovascular outcomes seeking correlations with COPD within the same cohort.

UPLIFT and TIOSPIR

Although not technically a cohort study, the Understanding Potential Long-Term Impact on Function with Tiotropium (UPLIFT) was a large interventional trial that included 5993 subjects with moderate-to-severe COPD.⁷ Patients worldwide were randomized to either the long-acting antimuscarinic agent tiotropium or placebo, in addition to their usual respiratory medications. The primary end point was rate of FEV₁ decline. Secondary end points included overall and respiratory-specific death. The addition of tiotropium conferred an improvement in FEV₁ decline, but, surprisingly, the rate of cardiac-specific death was also *reduced* in the tiotropium group⁸ (HR 0.86, 95% CI 0.75 to 0.99), despite similar smoking rates of ~30%. UPLIFT identified a subgroup of patients with COPD and cardiovascular disease (occult or known) who benefited from COPD-specific therapy, though pre-existing cardiovascular disease was not among the inclusion/exclusion criteria.

A similar study, Tiotropium Safety and Performance in Respimat (TIOSPIR), randomized subjects with COPD to inhaled tiotropium in different doses and different inhaler delivery devices on top of their usual non-anticholinergic medications.⁹ TIOSPIR included >17,000 subjects with GOLD 2–4 disease, and patients with stable cardiovascular disease were included. In addition to showing that tiotropium inhaled as a dry powder using the Handihaler or as a soft mist using the Respimat was equally effective in standard COPD outcomes, TIOSPIR found overall low rates of cardiac events (0.1–0.2% myocardial infarction (MI), 1.2–1.4% cardiac death). The authors found no evidence that one delivery device for tiotropium is safer than the other or was associated with a greater risk of major adverse cardiovascular events.

It is known that a substantial proportion of patients with COPD die from cardiovascular disease,¹⁰ an ‘overlap group’, and the UPLIFT and TIOSPIR trials suggest that this group benefits from COPD treatment. It is noteworthy, however, that some studies have not supported the concept that an ‘overlap group’ may derive cardiovascular benefit from COPD treatment.^{11–15} An early meta-analysis by Singh *et al*¹² suggested as much as a 52% increased risk of mortality associated with tiotropium mist inhaler use in patients with COPD, with another meta-analysis supporting a similar conclusion.¹³ However, the weight of the evidence, including the large randomized TIOSPIR trial that featured a prespecified subgroup analysis involving patients with underlying cardiovascular disease, supports that tiotropium is safe as the overall hazard of major adverse cardiovascular events, including death, was not increased.^{9,16} Postmarketing surveillance focused on cardiovascular events was recommended by the authors to validate their study findings.

Phenotyping and biomarkers

Thoracic CT scanning

One common feature in each of the three population studies COPDGene, ECLIPSE, and MESA-LUNG is the incorporation of CT scans to identify novel radiography-based ‘biomarkers’. Washko *et al*¹⁷ validated that CT scan-based measurements of airway wall attenuation are reproducible and correlate to the FEV₁/FVC ratio. This

study suggests that airway measurements by CT could be complementary to spirometry. A related study determined that the total number of small airways inversely correlated with the percent of emphysema, and that total airway count was predictive of BODE score (the prognostication metric calculated by assessing Body mass index, degree of airflow obstruction, degree of Dyspnea, and Exercise capacity).¹⁸

Building further on the relationship between airway size, caliber, and parenchymal changes, researchers established that the distensibility of medium-sized to large-sized airways is reduced in individuals with a predominantly emphysema phenotype versus an airway inflammatory phenotype on CT.¹⁹ When Martinez *et al*²⁰ assessed the correlation between radiological features of COPD, quality of life, and symptom measures, they discovered that patients with airway-limited disease had worse St George Respiratory Questionnaire (SGRQ) scores while those with more emphysema had increased (worse) BODE scores. Measures of air trapping, defined as low attenuation areas of <856 Hounsfield units, were additive to the value of airway measures alone in correlating with FEV₁ and FEV₁/FVC ratio.²¹ The presence of emphysema, separate from evidence of airflow limitation, was found to be associated with a lower total FEV₁ and worse functional status.²² ECLIPSE showed a higher risk of emphysema progression in women and active smokers. A similar risk related to gender and African-American ethnicity was identified by the COPDgene group.²³ The biomarkers surfactant protein D (SP-D) and soluble receptor for advanced glycation endproducts were more common in the progressive emphysema cohort.²⁴ However, correlation between COPD and cardiovascular disease outcomes was not the primary purpose of these studies.

Using MRI and CT scans, the MESA-LUNG and MESA-COPD investigators were the first to report that pulmonary microvascular changes are present in patients with mild, moderate, and severe COPD (defined by reduced FEV₁).²⁵ This study identified a decrease in the microvascular blood flow that was separate from the degree of emphysema present in those areas. The MESA-LUNG group also reported that CT evidence of pulmonary emphysema occurred in smokers with and without COPD, and that this emphysema was associated with symptoms if it was anatomically centrilobular or panlobular but not paraseptal.²⁶ MESA-LUNG investigators also commented on the relationship between emphysema and impaired left ventricular filling, concluding that pulmonary vein dimensions are reduced in patients with emphysema and COPD.²⁷

Serum biomarkers

Chronic persistent inflammation is generally thought to be a central feature of COPD, despite very little evidence that systemic anti-inflammatory therapy improves markers of inflammation. Assessment and discovery of distinct inflammatory patterns in COPD was a goal of all of the prospective cohort studies. Interestingly, Bowler *et al*²⁸ discovered that decreased levels of interleukin (IL)-16 were associated with emphysema²⁸ and may be related to the development of autoimmunity.

The hypothesis of systemic inflammation was most comprehensively explored by the ECLIPSE investigators. They

found that inflammation is not present in all patients with COPD, but when present, it appeared to be associated with poorer outcomes.²⁹ Additionally, they found that combining biomarkers as a composite score of inflammation, including C reactive protein (CRP), fibrinogen, and white cell count, was associated with more frequent exacerbations and comorbidities.³⁰ Fibrinogen was found to be elevated in 36% of patients with COPD as compared with 5% of control patients, and this has been identified as a candidate biomarker to identify patients at higher risk of frequent exacerbations, hospitalization, or mortality.³¹ Interestingly, fibrinogen is also known as a biomarker of cardiac disease.³²

Breath biomarkers

Breath biomarkers are an attractive and novel way to study COPD phenotypes as they are largely non-invasive and may complement existing biomarkers of disease. Until now, there have been efforts to use breath metabolites as a diagnostic matrix from patients with developing COPD^{33–35} and smokers at risk of COPD.³⁶ Studies of exhaled breath condensate (EBC)—the liquid formed from breath passed through a cold tube—identified lower fluid pH and higher hydrogen peroxide levels correlating with COPD.^{37–39} Other efforts have looked at EBC conductivity in emphysema,⁴⁰ EBC α -1-antitrypsin levels in acute COPD exacerbations,⁴¹ and fractional exhalation of nitric oxide in subjects with COPD.⁴² Although these studies show promise, differing biomarker collection techniques and analytic methods make standardization problematic, and larger scale studies and standardized procedures will surely advance the field. There are limited studies of exhaled biomarkers in patients with cardiovascular disease. Still, non-invasive and low-risk assessment tools that may add an important dimension to phenotyping COPD make breath analysis an exciting area of research.

COPD-associated comorbidities

Possibly as a consequence of systemic inflammation, patients with COPD are at a higher risk of developing associated diseases independent of smoking-induced airway disease. COPDgene researchers reported a relationship between COPD and cardiovascular disease. Matsuoka *et al*⁴³ showed that the cross-section of small pulmonary arteries correlates with the degree of aortic calcification. Another study reported that distal pruning of the pulmonary vasculature is a characteristic signature of smoking-related lung disease and associated with accelerated loss of lung tissue.⁴⁴ Researchers have established that seven common comorbid conditions are associated with COPD, including sleep apnea, stroke, coronary disease, peripheral vascular disease, osteoporosis, gastroesophageal reflux, and congestive heart failure (CHF).^{45–47} These associations are more pronounced among African-Americans.⁴⁷ Additionally, cardiovascular disease was independently associated with COPD.⁴⁸ The prevalence of venothromboembolic disease was higher in patients with COPD and comorbid conditions, and the overlap leads to worse exercise performance.⁴⁹ Finally, two separate investigations reported an increased association between COPD and diabetes mellitus.^{50 51} Similar findings were noted in the ECLIPSE cohort, where comorbid COPD and cardiovascular disease were associated with more

symptoms.³¹ Additionally, diabetes was identified as increasing the risk of poor clinical outcomes when associated with COPD. Depression was also identified as being more prevalent in COPD.⁵²

COPD immunobiology with a focus on vascular disease

COPD leads to anatomic distortion of normal airway architecture, resulting in a critical reduction in airway diameter and airflow limitation.⁵³ The major mechanisms thought responsible for airflow limitation include accumulated debris and mucus in the airway lumen, chronic bronchoconstriction, airway wall thickening, and increased external airway compression from a loss of elastic tissue. However, the rate of development of airflow limitation, that is, lung function loss, varies widely between patients with COPD. Factors such as quantity and quality of toxicant exposure (eg, tobacco smoke), innate and adaptive immune responses, and genetic and epigenetic elements that regulate airway inflammation and remodeling all contribute to the clinical progression in any single person. Clearly, the interplay between immune cells, toxicant exposure, and host background is complex and may evolve over the life of the patient with COPD.

Since COPD stems from abnormal lung and systemic inflammation, and advanced COPD is associated with comorbid vascular disease, there is considerable interest in understanding the immunological links between lung and vascular inflammation. It is known that COPD and coronary arterial disease (CAD) are connected,^{54–56} and the dominant theory is that shared risk factors (eg, smoking) elicit a chronic inflammatory response that affects both the lungs and vasculature.^{56–58} In fact, patients with COPD with elevated levels of systemic inflammatory markers such as CRP, fibrinogen, and leukocytes have increased rates of MI and CHF based on large cohort studies.⁵⁹ Efforts to unravel genetic links by comparing COPD-specific single nucleotide polymorphisms to carotid thickness and CAD are underway.⁶⁰ While at present a clear connection is not well established, it is imperative to understand concepts of shared cellular and molecular pathways such as oxidative stress, cell death, airway structural changes and impaired tissue repair underlying both chronic vascular conditions and COPD. The following sections discuss pulmonary structural and inflammatory cells in COPD with a focus on how these may relate to vascular inflammation (figure 1).

Immune cells, inflammation, and the lung-vascular connection

The gross insult by tobacco smoke to the respiratory tract is the result of repeated and prolonged exposure to a range of toxicants through inflammation and oxidative stress, or to individual toxicants through specific mechanisms.⁶¹ In COPD, damaged epithelial cells express high levels of inflammatory mediators (chemokine (C-X-C motif) ligand (CXCL)-8, IL-1- β , and granulocyte-macrophage colony-stimulating factor)⁶² and adhesion molecules (soluble intercellular adhesion molecule-1⁶² and E-selectin⁶³). This inflammatory response facilitates a continuous recruitment and activation of inflammatory cells from the blood. In addition, damaged lung epithelial cells have an altered ability to regulate normal immune functions such as pathogen binding,⁶⁴ antigen presentation, and tumor necrosis

factor- α (TNF- α) expression.^{65–68} In addition to its proinflammatory function, the airway epithelium is also responsible for maintaining immune homeostasis and protection against chronic inflammatory changes in the lung and the pulmonary vasculature. The protective function of airway epithelial cells has been attributed to the constitutive production of lung immune modulators called collectins: SP-A and SP-D. Although we currently do not have any direct evidence of a shared mechanism, SP-D-related immune regulatory pathways can be impaired in the development of atherosclerotic plaques⁶⁹ and increased levels of SP-D have been observed in heart failure⁷⁰ and carotid artery atherosclerosis.⁷¹ These data suggest that SP-D may be a biomarker or may play a putative role in coexistent lung and vascular disease.

Alveolar macrophages are the most abundant immune cell type in the lungs and airways. They function to clear inhaled particles, identify and destroy pathogens, and remove dead or dying cells in the distal air spaces. In COPD, however, the function of these cells is severely impaired⁷² despite increased numbers of macrophages in patients with COPD.^{73–74} In fact, macrophages, activated locally or recruited during inflammation, can account for many of the known features of COPD.^{74–75} Macrophages isolated from the lungs of patients with COPD exhibit reduced apoptosis and increased survival compared with those found in patients with normal lungs. Though this increased survival may be anti-inflammatory in the lung, damaged lung macrophages can produce IL-6,^{56–76} which in turn can potentiate coronary endothelial dysfunction.⁷⁷ Indeed, bone marrow-derived macrophages in the COPD lung differentiate into the highly proinflammatory M1 subtype and the anti-inflammatory M2 subtype; M1 macrophages have a well-accepted pathogenic role in atherosclerosis and CAD.

Normal, healthy lung parenchyma contains few if any neutrophils. In COPD, damaged epithelial cells, activated macrophages, and T cells (via CXCL-8, CXCL-1, and leukotriene B₄) cause direct migration of neutrophils toward the airways. Adhesion molecules expressed on endothelial and epithelial cells mediate neutrophil migration with the MAC1/ICAM1 interactions being the most crucial, and patients with COPD who smoke have increased surface expression of MAC1 on their neutrophils.⁷⁸ Neutrophils play a major role in COPD exacerbations elicited by air pollution, viral, and/or bacterial infections.^{79–81} Recruited neutrophils secrete a number of proinflammatory cytokines that elicit reactive oxygen species (ROS) formation, which further perpetuates neutrophil recruitment.⁸² Oxidative stress also causes elevated levels of cytokine and growth factor expression responsible for activating and preventing apoptosis of neutrophils. This effect can lead to either increased survival or necrotic death of these cells. An important feature of the COPD lung is an increased number of dead neutrophils due to necrotic cell death and a reduced ability of alveolar macrophages to perform their scavenger function. As with chronically activated macrophages, chronic neutrophil activity can lead to repeated endothelial exposure to cytotoxic agents (ie, ROS such as myeloperoxidase) and likely potentiate inflammatory changes, recurrent vasoconstriction, and cholesterol dysregulation.⁸³ In particular, neutrophil-derived ROS can potentiate elastin degradation which has been associated

GOLD Group	First Choice*	Alternate Choice
A FEV ₁ ≥ 50%** Low symptoms† <2 exacerbations/yr	Short-acting bronchodilators prn OR Long-acting bronchodilator (LABA or LAMA)	LAMA or LABA added
B FEV ₁ ≥ 50%** High symptoms† <2 exacerbations/yr	Long-acting bronchodilator (LABA or LAMA) OR LABA + LAMA	
C FEV ₁ < 50%** Low symptoms† ≥2 exacerbations/yr	LAMA OR LABA + LAMA	LABA + ICS
D FEV ₁ < 50%** High symptoms† ≥2 exacerbations/yr	LABA + LAMA OR LABA + LAMA + ICS	LAMA OR LABA + ICS OR LABA + LAMA + ICS + roflumilast OR LABA + LAMA + ICS + macrolides

Figure 1 Recommended therapy for stable COPD by GOLD category. Figure adapted from the recommendations of the GOLD¹ (<http://www.goldcopd.org>, accessed Jan 2017). *First choice therapy includes short-acting β -2-agonists or short-acting anticholinergic medications as needed for all categories. First choice therapy also includes the first entry followed by a clinical evaluation. If the patient still has symptoms, then moving to the second entry is advised. **FEV₁ impairment is FEV₁ ≥ 50% predicted for GOLD categories A and B, and FEV₁ < 50% predicted for categories C and D. †Symptoms based on the mMRC scale: 0—SOB with strenuous exertion, 1—SOB with hurrying on level ground or inclines, 2—SOB with normal walking on level ground >100 m, 3—SOB within 100 m, and 4—SOB with daily activities; mMRC < 2 = low symptoms and mMRC ≥ 2 = high symptoms. COPD, chronic obstructive pulmonary disease; CXCL, chemokine (C-X-C motif) ligand; FEV₁, forced expiratory volume in 1 s; GM-CSF, granulocyte-macrophage colony-stimulating factor; GOLD, Global Initiative for Chronic Obstructive Lung Disease; ICAM, intercellular Adhesion molecule; ICS, inhaled corticosteroid; LABA, long-acting β -2 agonist; LAMA, long-acting antimuscarinic; MAC, macrophage adhesion ligand; mMRC, modified Medical Research Council; SOB, shortness of breath; VCAM, vascular cell adhesion molecule.

with significant comorbid cardiac disease in patients with COPD.⁸⁴ Although the exact association between lung neutrophil activity and cardiovascular disease is not entirely clear, clinical evidence links circulating myeloperoxidase levels with adverse cardiac outcomes.^{85–87}

Lymphocyte accumulation in the pulmonary interstitium and peribronchial areas correlate with the severity of the symptoms of COPD and are considered to be part of the mechanism leading to exacerbation of symptoms brought on by air pollution or infections.⁷⁹ Lymphocytes organized in follicular structures with B lymphocyte-containing germinal centers surrounded by CD4+ T helper 1 (Th1) cells have been observed in clinically advanced cases of chronic bronchitis, while increases in the numbers of CD8+ cytotoxic Tc1 lymphocytes in the alveolar wall appear to be proportional to the severity of emphysema.⁸⁸ Th1 cells are CD4+ T cells that lead to interferon- γ secretion, and this, in turn, helps activate CD8+ cytotoxic Tc1 cells.⁸⁹ CD8+ T cells synthesize, store, and release cytokines and cytotoxic substances like TNF- α , granzyme B, and perforins, and their numbers inversely correlate with the FEV₁ of patients suffering from COPD.⁹⁰

Pulmonary endothelial cells, COPD, and vascular effects

Often described as the silent player in COPD pathogenesis, the pulmonary vasculature has been increasingly recognized as a major contributor to disease. Beyond their physiological function, endothelial cells also secrete a variety of proinflammatory molecules including cytokines, chemokines,

growth factors, and lipids relevant to COPD.⁹¹ Since COPD itself is a systemic inflammatory condition, both the systemic and pulmonary vasculature have enhanced expression of adhesion molecules (eg, vascular cell adhesion molecule-1) which further promote adherence of activated leukocytes to endothelial surfaces.⁷⁶ The pulmonary and airway vasculature also express vascular endothelial growth factor (VEGF) and various adhesion molecules important in the immune response that mediates the transmigration of neutrophils to the airways. As described above, the inflammatory milieu in COPD likely correlates with cardiovascular disease through, in part, endothelial dysfunction. However, a recent study by Chandra *et al*⁹² challenges this notion as the authors did not find a significant correlation between endothelial dysfunction and reduced lung function (FEV₁) in cohorts of patients with atherosclerotic disease. This study underscores the need to better define patients with COPD based on biological parameters other than lung function in order to truly understand the link between COPD and cardiovascular disease.

Alterations in the structure of the pulmonary vasculature in COPD contribute to the development of pulmonary arterial hypertension (PAH) which is associated with reduced survival in COPD and has a higher prevalence in more advanced disease.⁹³ The underlying dysfunction of the endothelial compartment in COPD leads to an imbalance between vasoconstrictive and vasodilatory mediators further contributing to the development of PAH. This imbalance is in part driven by cigarette smoke which also damages pulmonary endothelial cells via protease activity,

dysregulated apoptosis, and oxidative stress.⁹⁴ The development of alveolar destruction and emphysema is in part also due to this vasculopathy. Pulmonary capillary septal endothelial cell apoptosis and reduced local alveolar production of VEGF and its receptor VEGFR2 also contribute to the development of emphysema. Interestingly, in healthy smokers who quit smoking, pulmonary capillary apoptosis is reversible. However, in patients with COPD, this mechanism of endothelial cell apoptosis continues to be active despite smoking cessation further contributing to the development of progressive airflow obstruction.⁹⁵ This may explain in part the continued decline in lung function over many years in COPD despite smoking cessation.

COPD treatment: focusing on comorbid cardiac disease

Current COPD therapies

Several excellent reviews of the pharmacological treatment of COPD have been written.^{96–98} The GOLD guidelines (2017) use patient grouping (groups A–D) based on spirometry (FEV₁), frequency of exacerbations, and burden symptoms as assessed by symptom scores to guide treatment considerations. In addition to smoking cessation and vaccines, GOLD treatment guidelines use a step-up approach based on groups A–D with the goals to reduce symptoms with combination bronchodilators and to reduce risks, particularly acute exacerbations with anticholinergic bronchodilators, and, if indicated, inhaled corticosteroids or roflumilast (figure 2). Table 1 summarizes the currently available combination inhalers for maintenance therapy. No current therapies have been demonstrated to change the natural course of COPD except for smoking cessation.

Novel and investigative COPD therapies

Several approaches to new drug therapy for COPD are ongoing. These include novel agents that are dual phosphodiesterase (PDE3 and PDE4) inhibitors and other agents that are more specific PDE4 inhibitors. Some of these can potentially be delivered by inhalation.⁹⁸ Novel macrolide/fluoroketolide compounds appear to have better anti-inflammatory profiles than current macrolides and may be useful in treating COPD. Agents that are antagonists of the human C-X-C chemokine receptor (CXCR)2 receptor modulating neutrophil trafficking have potential in the treatment and prevention of COPD. The p38 mitogen-activated protein kinase inhibitors also have potential in COPD.⁹⁸

Agents that antagonize matrix metalloproteinases have the potential to inhibit the development of emphysema and small airway fibrosis in animal models but none have been effective in humans. Many new biologic therapies have potential use in the treatment of COPD including humanized monoclonal antibodies directed at IL-5 and IL-17 receptors. Phosphoinositide-3 kinase inhibitors, soluble epoxide hydrolase inhibitors and orally active, γ -selective retinoid agonists are new potential approaches to treating COPD.⁹⁸ Exciting new approaches to the treatment and prevention of COPD are on the horizon.

The statin drugs (statins) have garnered much interest as a potential therapy for COPD. Despite several large retrospective studies that suggest that statins have a benefit in preserving lung function and reducing mortality and morbidity in patients with COPD,⁹⁹ prospective studies have

failed to show an advantage in patients with COPD without a significant cardiovascular risk factor.¹⁰⁰ The STATCOPE clinical trial did not show that simvastatin reduced exacerbations in patients with moderate-to-severe COPD.¹⁰¹ However, smaller clinical trials with pravastatin did show benefit in patients with COPD. In two randomized clinical trials, pravastatin was associated with increased exercise time and reduced systemic inflammation in COPD,¹⁰² and in patients with COPD with pulmonary hypertension treatment with pravastatin increased functional capacity and exercise time, reduced systolic pulmonary pressures, and improved the BORG dyspnea score.¹⁰³ Based on these data in sum, statins cannot be recommended for the treatment of COPD, especially with the results of the STATCOPE trial. However, one limitation in the STATCOPE study is that it did not include patients with COPD with overt cardiovascular disease or those with significant cardiac risk factors. STATCOPE excluded the very group of patients with COPD who benefited from statin use in multiple observational studies.¹⁰⁴

Cardiac treatment in patients with COPD

The association of tobacco use and COPD is unequivocal and puts patients with COPD at a higher risk for cardiovascular comorbidities.¹⁰⁵ Patients with COPD are more likely to have cardiovascular disease than matched non-COPD populations (OR=2.46, 95% CI 2.02 to 3.00, $p<0.00001$).¹⁰⁶ This includes a 2–5 time increased risk for MI, cardiac dysrhythmia, CHF, disease of the pulmonary vasculature, and peripheral vascular diseases. Hypertension is also more common in patients with COPD (OR=1.33, 95% CI 1.13 to 1.56, $p=0.0007$).¹⁰⁶ Medications used to treat these cardiovascular comorbidities such as diuretics and β -blockers can have potential detrimental drug–disease interactions and effects in patients with COPD.

The treatment of hypertension in patients with COPD has been reviewed elsewhere.¹⁰⁷ Thiazide (hydrochlorothiazide, chlorthalidone) and loop (furosemide, bumetanide, torsemide) diuretics used in the treatment of hypertension and CHF can cause serious toxicity through urinary potassium losses. This can be exacerbated when diuretics are used with inhaled β -2-receptor agonists, which cause the movement of potassium into the cell. This combination can lead to severe hypokalemia. These drugs can also generate a volume-contraction metabolic alkalosis leading to a further suppression in ventilatory drive, thus resulting in worsening hypoxemia and hypercapnia. Alkalemia also increase the risk for cardiac arrhythmias further exacerbating cardiac disease.

ACE inhibitors are effective in the control of hypertension and the treatment of CHF in patients with COPD. However, since 5–20% of the patients on ACE inhibitors can develop a cough, they must be used with caution in COPD. Prior use of ACE inhibitors has been shown to reduce mortality in patients with COPD admitted with exacerbations.¹⁰⁷ Although ACE inhibitors have been suggested to improve skeletal muscle function in patients with COPD, a recent randomized controlled 3-month trial of the ACE inhibitor fosinopril in patients with COPD failed to show improvement in strength of the quadriceps or exercise performance.¹⁰⁸




Figure 2 Schematic showing the overlap between dysfunctional lung structural cells and inflammatory cells in COPD that may have a connection to cardiovascular disease. COPD, chronic obstructive pulmonary disease; IL, interleukin; SP-D, surfactant protein D; TNF, tumor necrosis factor; VEGF, vascular endothelial growth factor.

Amiodarone is a class III antiarrhythmic drug used to treat complex and life-threatening cardiac arrhythmias. The use of amiodarone is associated with significant pulmonary toxicity. In a large study of patients with atrial fibrillation, amiodarone use was associated with a nearly 40% increase in pulmonary toxicity in men compared with women (HR=1.37, 95% CI 1.19 to 1.57, $p<0.0001$) and more than a doubled risk in pulmonary toxicity was seen in patients with COPD (HR=2.53, 95% CI 2.21 to 2.89, $p<0.0001$).¹⁰⁹ Approximately 3.1% of patients with atrial fibrillation without pre-existing pulmonary disease were found to have pulmonary toxicity after 4 years of taking amiodarone compared with 5.9% ($p=0.015$) of those patients with pre-existing pulmonary disease in another study.¹¹⁰ Patients with CHF and COPD who were treated with amiodarone and survived at least 1 year had a significantly greater decrease in lung diffusion capacity (DLCO) compared with patients treated with placebo (2.05 vs 0.09 mL/min per mm Hg, $p=0.008$) but had no difference in survival free of cardiac deaths.¹¹¹ Taken together, these limited data suggest that the risk–benefit ratio must be considered before treating patients who have significant COPD with amiodarone and they need to be carefully monitored with chest imaging and DLCO measurements while on amiodarone.

As noted above, the risk of cardiovascular disease is increased in patients with COPD.¹⁰⁶ The β -blockers are used widely in the treatment of CHF, hypertension, atrial fibrillation, and MI. Non-selective β -blockers such as propranolol have been shown to have a negative effect on lung function (FEV₁, FVC, and FEV₁% predicted) as compared with β -1 selective receptor blockers like atenolol. This effect holds

true both at baseline and after albuterol inhalation in patients with COPD or asthma.^{112 113} Non-selective β -blocking agents should therefore be avoided in patients with COPD in favor of the more selective β -1-receptor blocker agents.

Use of labetalol, a non-selective β -blocker that also blocks α -1-receptor, did not affect FEV₁ or the mid-expiratory flow volumes in patients with COPD and hypertension 2 hours after the administration of the maximum labetalol dose.¹¹⁴ Another non-selective β -blocker/ α -1-receptor blocker, carvedilol, was studied in patients with CHF and COPD and compared with the selective β -1-blockers metoprolol and bisoprolol. A 6 min walk and left ventricular ejection fraction did not change with the three drugs. However, FEV₁ was lowest with carvedilol, better in metoprolol, and best in the patients treated with bisoprolol.¹¹⁵ In patients with CHF with COPD ($n=31$) or asthma ($n=12$), 3.2% of patients with COPD and 25% of patients with asthma developed wheezing after starting carvedilol.¹¹⁶ In contrast, actual improvement in peak expiratory flow rate of 17% ($p=0.04$) was seen in patients with COPD and 4% ($p=NS$) in patients with asthma 2 hours after starting carvedilol. The β and α adrenergic blocking agents should be used with caution in patients with COPD until more information is available.

In patients with COPD who had an MI, those discharged on β -blockers compared with those who did not had a lower all-cause mortality after adjusting for confounders (HR=0.87, 95% CI 0.64 to 0.95) during a follow-up period that was as long as 7.2 years.¹¹⁷ More impressive was the survival advantage seen in those patients with COPD discharged on a β -blocker after an MI and who also had CHF (HR=0.77, 95% CI 0.63 to 0.95).

Table 1 Combination drug inhalers used for maintenance treatment of COPD

Drug 1+drug 2	DI (μ g)	DF	Type DD	DD name	Name	
LABA+LAMA						
Indacaterol	Glycopyrronium	110/50	Once a day	DP	Ultibro Breezhaler	QVQ149
Vilanterol	Umeclidinium	25/62.5	Once a day	DP	Ellipta	Anoro
Olodaterol	Tiotropium	3.5/2.5	Once a day	SDM	Respimat	Stiolto
Formoterol	Acclidinium	12/400	two times a day	DP	Genuair	Duaklir
Formoterol	Glycopyrrolate	4.8/9	two times a day	MDI		Bevespi Aerosphere
SABA+SAMA						
Albuterol	Ipratropium	2.5/0.5 (mg)	Every 6 hours	Neb		DuoNeb
Albuterol	Ipratropium	0.1/0.33 (mg)	Every 6 hours	SDM	Respimat	Combivent
LABA +ICS						
Vilanterol	Fluticasone F	25/100	Once a day	DP	Ellipta	Breo
Formoterol	Budesonide	4.5/160 or 80	Two times a day	MDI		Symbicort
Formoterol	Budesonide	6612/100,200,400	Two times a day	DP	Turbuhaler	Symbicort
Formoterol	Mometasone	5/100 or 200	Two times a day	MDI		Dulera ^A
Salmeterol	Fluticasone P	50/100,250,500	Two times a day	DP	Diskus	Advair
Salmeterol	Fluticasone P	21/45,115,230	Two times a day	MDI		Advair HFA ^A

^A, approved for asthma indication only, all others approved for COPD or COPD+asthma; COPD, chronic obstructive pulmonary disease; DF, dose frequency; DI, dose per inhalation; DP, dry powder; F, furoate; ICS, inhaled corticosteroid; LABA, long-acting β -2 agonist; LAMA, long-acting muscarinic antagonist; MDI, metered dose inhaler; Neb, nebulization; P, propionate; SABA, short-acting β -2 agonist; SAMA, short-acting muscarinic antagonist; SDM, spring driven mist; Type DD, type of delivery device.

Meta-analysis of the use of selective β -1-receptor blockers for hypertension, CHF, and coronary artery disease and during the perioperative period in patients with COPD concluded that they did not produce adverse respiratory effects.¹¹⁸ However, a large prospective cohort observational trial showed that both cardioselective and non-cardioselective β -blockers in patients without lung disease were associated with significant reductions in FEV₁ measures over a mean of 6.1 \pm 0.5 years. The use of selective β -1-blockers resulted in less reduction in FEV₁ (–118 mL, 95% CI –157 to –78, p <0.001) than the reduction seen with the use of non-cardioselective β -blockers (–198 mL, 95% CI –301 to –96, p <0.001).¹¹⁹ When patients with COPD, asthma, and CHF were included, the same trends held.

In a clinical trial where patients with COPD and CHF were randomized to either the selective β -1-blocker bisoprolol or the non-selective β -blocker/ α -1-blocker carvedilol, both agents reduced the heart rate and had no effect on the N-terminal pro brain natriuretic peptide. Bisoprolol, but not carvedilol, significantly increased FEV₁ by 127 mL.¹²⁰ Another randomized triple cross-over trial evaluated carvedilol, metoprolol, and bisoprolol in patients with CHF and found that in those patients with COPD, bisoprolol had the highest and carvedilol the lowest FEV₁ measurements.¹¹⁵ However, bisoprolol use is also associated with worsening dynamic hyperinflation compared with placebo in patients with moderate-to-severe COPD without reducing the duration of exercise.¹²¹ Conversely, the rate of CHF and/or COPD exacerbations were higher in those patients treated with carvedilol as compared with bisoprolol.¹²²

Beyond lung function, β -blockers have been associated with important hard outcomes such as mortality. A mortality advantage was seen with the use of bisoprolol, but not carvedilol or metoprolol, in patients with COPD and CHF.¹²³ Another study demonstrated reduced mortality

rates in patients with COPD with CHF on bisoprolol or carvedilol (HR=0.41, 95% CI 0.17 to 0.99, p =0.047). In a large Scottish retrospective cohort study of β -blockers with a mean follow-up of 4.35 years, there was a 22% reduction in overall mortality in patients with COPD taking β -blockers.¹²⁴

Two large trials have demonstrated significant reductions in COPD exacerbations regardless of the severity of airflow obstruction when the patients are on β -blockers.^{125–126} A trial of 520 patients with COPD undergoing lung resection found that the use of perioperative β -blockers compared with not using them did not change the rate of postoperative COPD exacerbations (5.4% vs 6.3%).¹²⁷ Selective β -1-receptor blockers appear to have an advantage over non-selective β -blockers in patients with COPD with CHF, hypertension and MIs, but the advantages have been small and not always consistent.

SUMMARY

COPD is the third most common cause of death worldwide. The definition of COPD is evolving, due to complex disease mechanisms, clinical heterogeneity, and variable immune response to inhaled toxicants and environmental pollutants. Large cohort studies are important to help define COPD phenotypes and identify useful biomarkers, and these studies give rise to important and testable clinical questions such as how patients with certain radiological features respond to therapeutic interventions. As our understanding of COPD immunobiology improves, we may better identify specific and effective immune-modulating therapies at various stages of COPD, including monoclonal antibodies in the current age of biologics and precision medicine. The recognition that COPD often coexists with cardiovascular disease underscores the link between these disorders. Therapies directed at both COPD and heart disease seem to confer benefit beyond treating each separately, and the future of COPD research and treatment approaches needs to bear this in mind.

Acknowledgements The authors graciously acknowledge the following funding agencies for this work: NIH-NHLBI K23HL127185 (MS); NIH-NIAID R21HL11612 (AH); NIH-NHLBI K08HL114882 (AAZ).

Funding National Heart, Lung, and Blood Institute, 10.13039/100000050, K08HL114882 (AAZ), K23HL127185 (MS), National Institute of Allergy and Infectious Diseases, 10.13039/100000060, R21HL11612 (AH).

Competing interests None declared.

Provenance and peer review Commissioned; externally peer reviewed.

REFERENCES

- 2017 GfCOLDG. The Global Strategy for the Diagnosis, Management, and Prevention of COPD. 2017. <http://goldcopd.org/>
- Wan ES, Cho MH, Boutaoui N, *et al.* Genome-wide association analysis of body mass in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2011;45:304–10.
- Rennard SI, Locantore N, Delafont B, *et al.* Identification of five chronic obstructive pulmonary disease subgroups with different prognoses in the ECLIPSE cohort using cluster analysis. *Ann Am Thorac Soc* 2015;12:303–12.
- Rennard SI. The promise of observational studies (ECLIPSE, SPIROMICS, and COPDGene) in achieving the goal of personalized treatment of chronic obstructive pulmonary disease. *Semin Respir Crit Care Med* 2015;36:478–90.
- Yoneyama K, Donekal S, Venkatesh BA, *et al.* Natural history of myocardial function in an adult human population: serial longitudinal observations from MESA. *JACC Cardiovasc Imaging* 2016;9:1164–73.
- Barr RG, Ahmed FS, Carr JJ, *et al.* Subclinical atherosclerosis, airflow obstruction and emphysema: the MESA Lung Study. *Eur Respir J* 2012;39:846–54.
- Tashkin DP, Celli B, Senn S, *et al.* A 4-year trial of tiotropium in chronic obstructive pulmonary disease. *N Engl J Med* 2008;359:1543–54.
- Celli B, Decramer M, Kesten S, *et al.* Mortality in the 4-year trial of tiotropium (UPLIFT) in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2009;180:948–55.
- Wise RA, Anzueto A, Cotton D, *et al.* Tiotropium Respimat inhaler and the risk of death in COPD. *N Engl J Med* 2013;369:1491–501.
- Lange P, Marott JL, Vestbo J, *et al.* Prediction of the clinical course of chronic obstructive pulmonary disease, using the new GOLD classification: a study of the general population. *Am J Respir Crit Care Med* 2012;186:975–81.
- Gershon A, Croxford R, Calzavara A, *et al.* Cardiovascular safety of inhaled long-acting bronchodilators in individuals with chronic obstructive pulmonary disease. *JAMA Intern Med* 2013;173:1175–85.
- Singh S, Loke YK, Enright PL, *et al.* Mortality associated with tiotropium mist inhaler in patients with chronic obstructive pulmonary disease: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2011;342:d3215.
- Dong YH, Lin HH, Shau WY, *et al.* Comparative safety of inhaled medications in patients with chronic obstructive pulmonary disease: systematic review and mixed treatment comparison meta-analysis of randomised controlled trials. *Thorax* 2013;68:48–56.
- Karner C, Chong J, Poole P. Tiotropium versus placebo for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2014;(7):CD009285.
- Verhamme KM, Sturkenboom MC, Brusselle GG. Use of tiotropium Respimat versus HandiHaler and mortality in patients with COPD. *Eur Respir J* 2014;43:1818–19.
- Tan CK, Say GQ, Geake JB. Long-term safety of tiotropium delivered by Respimat(R) SoftMist Inhaler: patient selection and special considerations. *Ther Clin Risk Manag* 2016;12:1433–44.
- Washko GR, Dransfield MT, Estepar RS, *et al.* Airway wall attenuation: a biomarker of airway disease in subjects with COPD. *J Appl Physiol* 2009;107:185–91.
- Diaz AA, Valim C, Yamashiro T, *et al.* Airway count and emphysema assessed by chest CT imaging predicts clinical outcome in smokers. *Chest* 2010;138:880–7.
- Diaz AA, Come CE, Ross JC, *et al.* Association between airway caliber changes with lung inflation and emphysema assessed by volumetric CT scan in subjects with COPD. *Chest* 2012;141:736–44.
- Martinez CH, Chen YH, Westgate PM, *et al.* Relationship between quantitative CT metrics and health status and BODE in chronic obstructive pulmonary disease. *Thorax* 2012;67:399–406.
- Schroeder JD, McKenzie AS, Zach JA, *et al.* Relationships between airflow obstruction and quantitative CT measurements of emphysema, air trapping, and airways in subjects with and without chronic obstructive pulmonary disease. *AJR Am J Roentgenol* 2013;201:W460–70.
- Castaldi PJ, San Jose Estepar R, Mendoza CS, *et al.* Distinct quantitative computed tomography emphysema patterns are associated with physiology and function in smokers. *Am J Respir Crit Care Med* 2013;188:1083–90.
- Foreman MG, Zhang L, Murphy J, *et al.* Early-onset chronic obstructive pulmonary disease is associated with female sex, maternal factors, and African American race in the COPDGene Study. *Am J Respir Crit Care Med* 2011;184:414–20.
- Coxson HO, Dirksen A, Edwards LD, *et al.* The presence and progression of emphysema in COPD as determined by CT scanning and biomarker expression: a prospective analysis from the ECLIPSE study. *Lancet Respir Med* 2013;1:129–36.
- Hueper K, Vogel-Claussen J, Parikh MA, *et al.* Pulmonary microvascular blood flow in mild chronic obstructive pulmonary disease and emphysema. The MESA COPD Study. *Am J Respir Crit Care Med* 2015;192:570–80.
- Smith BM, Austin JH, Newell JD Jr, *et al.* Pulmonary emphysema subtypes on computed tomography: the MESA COPD study. *Am J Med* 2014;127:94.e7–23.
- Smith BM, Prince MR, Hoffman EA, *et al.* Impaired left ventricular filling in COPD and emphysema: is it the heart or the lungs? The Multi-Ethnic Study of Atherosclerosis COPD Study. *Chest* 2013;144:1143–51.
- Bowler RP, Bahr TM, Hughes G, *et al.* Integrative omics approach identifies interleukin-16 as a biomarker of emphysema. *OMICS* 2013;17:619–26.
- Agusti A, Edwards LD, Rennard SI, *et al.* Persistent systemic inflammation is associated with poor clinical outcomes in COPD: a novel phenotype. *PLoS ONE* 2012;7:e37483.
- Thomsen M, Ingebrigtsen TS, Marott JL, *et al.* Inflammatory biomarkers and exacerbations in chronic obstructive pulmonary disease. *JAMA* 2013;309:2353–61.
- Miller J, Edwards LD, Agusti A, *et al.* Comorbidity, systemic inflammation and outcomes in the ECLIPSE cohort. *Respir Med* 2013;107:1376–84.
- Kaptoge S, Di Angelantonio E, Pennells L, *et al.* Emerging Risk Factors Collaboration. C-reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med* 2012;367:1310–20.
- Besa V, Teschler H, Kurth I, *et al.* Exhaled volatile organic compounds discriminate patients with chronic obstructive pulmonary disease from healthy subjects. *Int J Chron Obstruct Pulmon Dis* 2015;10:399–406.
- Basanta M, Ibrahim B, Dockry R, *et al.* Exhaled volatile organic compounds for phenotyping chronic obstructive pulmonary disease: a cross-sectional study. *Respir Res* 2012;13:72.
- Schivo M, Seichter F, Aksenov AA, *et al.* A mobile instrumentation platform to distinguish airway disorders. *J Breath Res* 2013;7:017113.
- Malerba M, Montuschi P. Non-invasive biomarkers of lung inflammation in smoking subjects. *Curr Med Chem* 2012;19:187–96.
- Murata K, Fujimoto K, Kitaguchi Y, *et al.* Hydrogen peroxide content and pH of expired breath condensate from patients with asthma and COPD. *COPD* 2014;11:81–7.
- Inonu H, Doruk S, Sahin S, *et al.* Oxidative stress levels in exhaled breath condensate associated with COPD and smoking. *Respir Care* 2012;57:413–19.
- MacNee W, Rennard SI, Hunt JF, *et al.* Evaluation of exhaled breath condensate pH as a biomarker for COPD. *Respir Med* 2011;105:1037–45.
- Stolk J, Fumagalli M, Viglio S, *et al.* Conductivity in exhaled breath condensate from subjects with emphysema and type ZZ alpha-1-antitrypsin deficiency. *COPD* 2015;12(Suppl 1):32–5.
- Koczulla AR, Noeske S, Herr C, *et al.* Alpha-1 antitrypsin is elevated in exhaled breath condensate and serum in exacerbated COPD patients. *Respir Med* 2012;106:120–6.
- Malerba M, Radaeli A, Olivini A, *et al.* Exhaled nitric oxide as a biomarker in COPD and related comorbidities. *Biomed Res Int* 2014;2014:271918.
- Matsuoka S, Yamashiro T, Diaz A, *et al.* The relationship between small pulmonary vascular alteration and aortic atherosclerosis in chronic obstructive pulmonary disease: quantitative CT analysis. *Acad Radiol* 2011;18:40–6.
- Estepar RS, Kinney GL, Black-Shinn JL, *et al.* Computed tomographic measures of pulmonary vascular morphology in smokers and their clinical implications. *Am J Respir Crit Care Med* 2013;188:231–9.
- Laforest L, Roche N, Devouassoux G, *et al.* Frequency of comorbidities in chronic obstructive pulmonary disease, and impact on all-cause mortality: a population-based cohort study. *Respir Med* 2016;117:33–9.
- Decramer M, Janssens W. Chronic obstructive pulmonary disease and comorbidities. *Lancet Respir Med* 2013;1:73–83.

- 47 Putcha N, Han MK, Martinez CH, *et al.* Comorbidities of COPD have a major impact on clinical outcomes, particularly in African Americans. *COPD* 2014;1:105–14.
- 48 Black-Shinn JL, Kinney GL, Wise AL, *et al.* Cardiovascular disease is associated with COPD severity and reduced functional status and quality of life. *COPD* 2014;11:546–51.
- 49 Kim V, Goel N, Gangar J, *et al.* Risk factors for venous thromboembolism in chronic obstructive pulmonary disease. *COPD* 2014;1:239–49.
- 50 Kinney GL, Black-Shinn JL, Wan ES, *et al.* Pulmonary function reduction in diabetes with and without chronic obstructive pulmonary disease. *Diabetes Care* 2014;37:389–95.
- 51 Hersh CP, Make BJ, Lynch DA, *et al.* Non-emphysematous chronic obstructive pulmonary disease is associated with diabetes mellitus. *BMC Pulm Med* 2014;14:164.
- 52 Hanania NA, Mullerova H, Locantore NW, *et al.* Determinants of depression in the ECLIPSE chronic obstructive pulmonary disease cohort. *Am J Respir Crit Care Med* 2011;183:604–11.
- 53 Ge MQ, Kokalari B, Flayer CH, *et al.* Cutting edge: role of NK cells and surfactant protein D in dendritic cell lymph node homing: effects of ozone exposure. *J Immunol* 2016;196:553–7.
- 54 Man SF, Leipsic JA, Man JP, *et al.* Is atherosclerotic heart disease in COPD a distinct phenotype? *Chest* 2011;140:569–71.
- 55 Williams MC, Murchison JT, Edwards LD, *et al.* Coronary artery calcification is increased in patients with COPD and associated with increased morbidity and mortality. *Thorax* 2014;69:718–23.
- 56 Roversi S, Roversi P, Spadafora G, *et al.* Coronary artery disease concomitant with chronic obstructive pulmonary disease. *Eur J Clin Invest* 2014;44:93–102.
- 57 Man SF, Van Eeden S, Sin DD. Vascular risk in chronic obstructive pulmonary disease: role of inflammation and other mediators. *Can J Cardiol* 2012;28:653–61.
- 58 Boschetto P, Beghe B, Fabbri LM, *et al.* Link between chronic obstructive pulmonary disease and coronary artery disease: implication for clinical practice. *Respirology* 2012;17:422–31.
- 59 Thomsen M, Dahl M, Lange P, *et al.* Inflammatory biomarkers and comorbidities in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2012;186:982–8.
- 60 Sabater-Lleal M, Malarstig A, Folkersen L, *et al.* Common genetic determinants of lung function, subclinical atherosclerosis and risk of coronary artery disease. *PLoS ONE* 2014;9:e104082.
- 61 Shepperd CJ, Newland N, Eldridge A, *et al.* Changes in levels of biomarkers of exposure and biological effect in a controlled study of smokers switched from conventional cigarettes to reduced-toxicant-prototype cigarettes. *Regul Toxicol Pharmacol* 2015;72:273–91.
- 62 Hellermann GR, Nagy SB, Kong X, *et al.* Mechanism of cigarette smoke condensate-induced acute inflammatory response in human bronchial epithelial cells. *Respir Res* 2002;3:22.
- 63 Di Stefano A, Maestrelli P, Roggeri A, *et al.* Upregulation of adhesion molecules in the bronchial mucosa of subjects with chronic obstructive bronchitis. *Am J Respir Crit Care Med* 1994;149:803–10.
- 64 Rosseau S, Guenther A, Seeger W, *et al.* Phagocytosis of viable *Candida albicans* by alveolar macrophages: lack of opsonin function of surfactant protein A. *J Infect Dis* 1997;175:421–8.
- 65 Brinker KG, Martin E, Borron P, *et al.* Surfactant protein D enhances bacterial antigen presentation by bone marrow-derived dendritic cells. *Am J Physiol Lung Cell Mol Physiol* 2001;281:L1453–63.
- 66 Brinker KG, Garner H, Wright JR. Surfactant protein A modulates the differentiation of murine bone marrow-derived dendritic cells. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L232–41.
- 67 Hansen S, Lo B, Evans K, *et al.* Surfactant protein D augments bacterial association but attenuates major histocompatibility complex class II presentation of bacterial antigens. *Am J Respir Cell Mol Biol* 2007;36:94–102.
- 68 Hortobagyi L, Kierstein S, Krytska K, *et al.* Surfactant protein D inhibits TNF- α production by macrophages and dendritic cells in mice. *J Allergy Clin Immunol* 2008;122:521–8.
- 69 Barrow AD, Palarasah Y, Bugatti M, *et al.* OSCAR is a receptor for surfactant protein D that activates TNF- α release from human CCR2+ inflammatory monocytes. *J Immunol* 2015;194:3317–26.
- 70 Gargiulo P, Banfi C, Ghilardi S, *et al.* Surfactant-derived proteins as markers of alveolar membrane damage in heart failure. *PLoS ONE* 2014;9:e115030.
- 71 Hu F, Zhong Q, Gong J, *et al.* Serum surfactant protein D is associated with atherosclerosis of the carotid artery in patients on maintenance hemodialysis. *Clin Lab* 2016;62:97–104.
- 72 Hodge S, Hodge G, Scicchitano R, *et al.* Alveolar macrophages from subjects with chronic obstructive pulmonary disease are deficient in their ability to phagocytose apoptotic airway epithelial cells. *Immunol Cell Biol* 2003;81:289–96.
- 73 Finkelstein R, Fraser RS, Ghezzi H, *et al.* Alveolar inflammation and its relation to emphysema in smokers. *Am J Respir Crit Care Med* 1995;152:1666–72.
- 74 Barnes PJ. Alveolar macrophages as orchestrators of COPD. *COPD* 2004;1:59–70.
- 75 Barnes PJ. Alveolar macrophages in chronic obstructive pulmonary disease (COPD). *Cell Mol Biol* 2004;50 Online Pub:OL627–37.
- 76 van Eeden SF, Sin DD. Chronic obstructive pulmonary disease: a chronic systemic inflammatory disease. *Respiration* 2008;75:224–38.
- 77 Zhang J, Alcaide P, Liu L, *et al.* Regulation of endothelial cell adhesion molecule expression by mast cells, macrophages, and neutrophils. *PLoS ONE* 2011;6:e14525.
- 78 Domagala-Kulawik J. Effects of cigarette smoke on the lung and systemic immunity. *J Physiol Pharmacol* 2008;59(Suppl 6):19–34.
- 79 Papi A, Luppi F, Franco F, *et al.* Pathophysiology of exacerbations of chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2006;3:245–51.
- 80 Tsoumakidou M, Siafakas NM. Novel insights into the aetiology and pathophysiology of increased airway inflammation during COPD exacerbations. *Respir Res* 2006;7:80.
- 81 Celli BR, Barnes PJ. Exacerbations of chronic obstructive pulmonary disease. *Eur Respir J* 2007;29:1224–38.
- 82 Rahman I. Oxidative stress in pathogenesis of chronic obstructive pulmonary disease: cellular and molecular mechanisms. *Cell Biochem Biophys* 2005;43:167–88.
- 83 King PT. Inflammation in chronic obstructive pulmonary disease and its role in cardiovascular disease and lung cancer. *Clin Transl Med* 2015;4:68.
- 84 Rabinovich RA, Miller BE, Wrobel K, *et al.* Circulating desmosine levels do not predict emphysema progression but are associated with cardiovascular risk and mortality in COPD. *Eur Respir J* 2016;47:1365–73.
- 85 Tang WH, Wu Y, Nicholls SJ, *et al.* Plasma myeloperoxidase predicts incident cardiovascular risks in stable patients undergoing medical management for coronary artery disease. *Clin Chem* 2011;57:33–9.
- 86 Meuwese MC, Stroes ES, Hazen SL, *et al.* Serum myeloperoxidase levels are associated with the future risk of coronary artery disease in apparently healthy individuals: the EPIC-Norfolk Prospective Population Study. *J Am Coll Cardiol* 2007;50:159–65.
- 87 Brennan ML, Penn MS, Van Lente F, *et al.* Prognostic value of myeloperoxidase in patients with chest pain. *N Engl J Med* 2003;349:1595–604.
- 88 Hogg JC, Chu F, Utokaparch S, *et al.* The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350:2645–53.
- 89 Barnes PJ, Rennard SI. Pathophysiology of COPD. In: Barnes PJ, Drazen JM, Rennard SI, Thomson NC, eds. *Asthma and COPD: basic mechanisms and clinical management*. 2nd edn. Elsevier, 2009.
- 90 Freeman CM, Han MK, Martinez FJ, *et al.* Cytotoxic potential of lung CD8(+) T cells increases with chronic obstructive pulmonary disease severity and with in vitro stimulation by IL-18 or IL-15. *J Immunol* 2010;184:6504–13.
- 91 Barnes PJ. Inflammatory mechanisms in patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol* 2016;138:16–27.
- 92 Chandra D, Gupta A, Strollo PJ Jr, *et al.* Airflow limitation and endothelial dysfunction. Unrelated and independent predictors of atherosclerosis. *Am J Respir Crit Care Med* 2016;194:38–47.
- 93 Blanco I, Piccirilli L, Barbera JA. Pulmonary vasculature in COPD: the silent component. *Respirology* 2016;21:984–94.
- 94 Voelkel NF, Cool CD. Pulmonary vascular involvement in chronic obstructive pulmonary disease. *Eur Respir J Suppl* 2003;46:28S–32S.
- 95 Strulovici-Barel Y, Staudt MR, Krause A, *et al.* Persistence of circulating endothelial microparticles in COPD despite smoking cessation. *Thorax* 2016. doi: 10.1136/thoraxjnl-2015-208274 [Epub ahead of print: 26 Jul 2016].
- 96 Albertson TE, Schivo M, Zeki AA, *et al.* The pharmacological approach to the elderly COPD patient. *Drugs Aging* 2013;30:479–502.
- 97 Bateman ED, Mahler DA, Vogelmeier CF, *et al.* Recent advances in COPD disease management with fixed-dose long-acting combination therapies. *Expert Rev Respir Med* 2014;8:357–79.
- 98 Barjaktarevic IZ, Arredondo AF, Cooper CB. Positioning new pharmacotherapies for COPD. *Int J Chron Obstruct Pulmon Dis* 2015;10:1427–42.

- 99 Dobler CC, Wong KK, Marks GB. Associations between statins and COPD: a systematic review. *BMC Pulm Med* 2009;9:32.
- 100 Howard ML, Vincent AH. Statin effects on exacerbation rates, mortality, and inflammatory markers in patients with chronic obstructive pulmonary disease: a review of prospective studies. *Pharmacotherapy* 2016;36:536–47.
- 101 Criner GJ, Connett JE, Aaron SD, *et al.* Simvastatin for the prevention of exacerbations in moderate-to-severe COPD. *N Engl J Med* 2014;370:2201–10.
- 102 Lee TM, Lin MS, Chang NC. Usefulness of C-reactive protein and interleukin-6 as predictors of outcomes in patients with chronic obstructive pulmonary disease receiving pravastatin. *Am J Cardiol* 2008;101:530–5.
- 103 Lee TM, Chen CC, Shen HN, *et al.* Effects of pravastatin on functional capacity in patients with chronic obstructive pulmonary disease and pulmonary hypertension. *Clin Sci* 2009;116:497–505.
- 104 Young RP, Hopkins RJ, Agusti A. Statins as adjunct therapy in COPD: how do we cope after STATCOPE? *Thorax* 2014;69:891–4.
- 105 Brown JP, Martinez CH. Chronic obstructive pulmonary disease comorbidities. *Curr Opin Pulm Med* 2016;22:113–18.
- 106 Chen W, Thomas J, Sadatsafavi M, *et al.* Risk of cardiovascular comorbidity in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Lancet Respir Med* 2015;3:631–9.
- 107 Chandy D, Aronow WS, Banach M. Current perspectives on treatment of hypertensive patients with chronic obstructive pulmonary disease. *Integr Blood Press Control* 2013;6:101–9.
- 108 Shrikrishna D, Tanner RJ, Lee JY, *et al.* A randomized controlled trial of angiotensin-converting enzyme inhibition for skeletal muscle dysfunction in COPD. *Chest* 2014;146:932–40.
- 109 Jackevicius CA, Tom A, Essebag V, *et al.* Population-level incidence and risk factors for pulmonary toxicity associated with amiodarone. *Am J Cardiol* 2011;108:705–10.
- 110 Olshansky B, Sami M, Rubin A, *et al.* Use of amiodarone for atrial fibrillation in patients with preexisting pulmonary disease in the AFFIRM study. *Am J Cardiol* 2005;95:404–5.
- 111 Singh SN, Fisher SG, Deedwania PC, *et al.* Pulmonary effect of amiodarone in patients with heart failure. The Congestive Heart Failure-Survival Trial of Antiarrhythmic Therapy (CHF-STAT) Investigators (Veterans Affairs Cooperative Study No. 320). *J Am Coll Cardiol* 1997;30:514–17.
- 112 Doshan HD, Rosenthal RR, Brown R, *et al.* Celiprolol, atenolol and propranolol: a comparison of pulmonary effects in asthmatic patients. *J Cardiovasc Pharmacol* 1986;8(Suppl 4):S105–8.
- 113 Fogari R, Zoppi A, Tettamanti F, *et al.* Comparative effects of celiprolol, propranolol, oxprenolol, and atenolol on respiratory function in hypertensive patients with chronic obstructive lung disease. *Cardiovasc Drugs Ther* 1990;4:1145–9.
- 114 George RB, Manocha K, Burford JG, *et al.* Effects of labetalol in hypertensive patients with chronic obstructive pulmonary disease. *Chest* 1983;83:457–60.
- 115 Jabbour A, Macdonald PS, Keogh AM, *et al.* Differences between beta-blockers in patients with chronic heart failure and chronic obstructive pulmonary disease: a randomized crossover trial. *J Am Coll Cardiol* 2010;55:1780–7.
- 116 Kotlyar E, Keogh AM, Macdonald PS, *et al.* Tolerability of carvedilol in patients with heart failure and concomitant chronic obstructive pulmonary disease or asthma. *J Heart Lung Transplant* 2002;21:1290–5.
- 117 Andell P, Erlinge D, Smith JG, *et al.* beta-blocker use and mortality in COPD patients after myocardial infarction: a Swedish nationwide observational study. *J Am Heart Assoc* 2015;4:pil: e001611.
- 118 Salpeter S, Ormiston T, Salpeter E. Cardioselective beta-blockers for chronic obstructive pulmonary disease. *Cochrane Database Sys Rev* 2005;(4): CD003566.
- 119 Loth DW, Brusselle GG, Lahousse L, *et al.* beta-adrenoceptor blockers and pulmonary function in the general population: the Rotterdam Study. *Br J Clin Pharmacol* 2014;77:190–200.
- 120 Lainscak M, Podbregar M, Kovacic D, *et al.* Differences between bisoprolol and carvedilol in patients with chronic heart failure and chronic obstructive pulmonary disease: a randomized trial. *Respir Med* 2011;105:S44–S9.
- 121 Mainguy V, Girard D, Maltais F, *et al.* Effect of bisoprolol on respiratory function and exercise capacity in chronic obstructive pulmonary disease. *Am J Cardiol* 2012;110:258–63.
- 122 Kubota Y, Asai K, Furuse E, *et al.* Impact of beta-blocker selectivity on long-term outcomes in congestive heart failure patients with chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis* 2015;10:515–23.
- 123 Su VY, Chang YS, Hu YW, *et al.* Carvedilol, bisoprolol, and metoprolol use in patients with coexistent heart failure and chronic obstructive pulmonary disease. *Medicine (Baltimore)* 2016;95:e2427.
- 124 Short PM, Lipworth SI, Elder DH, *et al.* Effect of beta blockers in treatment of chronic obstructive pulmonary disease: a retrospective cohort study. *BMJ* 2011;342:d2549.
- 125 Bhatt SP, Wells JM, Kinney GL, *et al.* beta-Blockers are associated with a reduction in COPD exacerbations. *Thorax* 2016;71:8–14.
- 126 Farland MZ, Peters CJ, Williams JD, *et al.* beta-Blocker use and incidence of chronic obstructive pulmonary disease exacerbations. *Ann Pharmacother* 2013;47:651–6.
- 127 Kamath A, Stover DE, Hemdan A, *et al.* Effect of perioperative beta-blockers on pulmonary complications among patients with chronic obstructive pulmonary disease undergoing lung resection surgery. *Lung Cancer Int* 2015;2015:204826.