



Evaluation of mesenteric artery disease in patients with severe aortic valve stenosis

Aysegul Idil Soylu ,¹ Ufuk Avcioglu,² Fatih Uzunkaya,¹ Korhan Soylu ³

¹Department of Radiology, Ondokuz Mayıs University, Faculty of Medicine, Samsun, Turkey

²Department of Gastroenterology, Ondokuz Mayıs University, Faculty of Medicine, Samsun, Turkey

³Department of Cardiology, Faculty of Medicine, Ondokuz Mayıs University, Samsun, Turkey

Correspondence to

Dr Aysegul Idil Soylu, Radiology, Ondokuz Mayıs University Faculty of Medicine, Samsun 55200, Turkey; a.isoylu@gmail.com

Accepted 3 November 2020
Published Online First
15 January 2021

ABSTRACT

The aim of this study is to evaluate the mesenteric artery stenosis (MAS) in routinely performed CT angiography (CTA) of patients with severe aortic stenosis (AS) planned for transcatheter aortic valve implantation (TAVI) before the procedure. Patients with AS (AS group) who routinely underwent CTA before the TAVI procedure due to severe AS and patients who had CTA for other indications (control group) were retrospectively and sequentially scanned. The demographic characteristics of the patients in both groups were similar. Calcification and stenosis in the mesenteric arteries were recorded according to the localization of celiac truncus, superior mesenteric artery (SMA) and inferior mesenteric artery (IMA). Class 0–3 classification was used for calcification score. Stenoses with a stenosis degree $\geq 50\%$ were considered as significant. A total of 184 patients, 73 patients with severe AS and 111 control groups, were included in the study. SMA and IMA calcification scores of patients with AS were significantly higher than the control group ($p=0.035$ for SMA and $p=0.020$ for IMA). In addition, the rate of patients with significant MAS in at least 1 artery (45.2% vs 22.5%, $p=0.001$) and the rate of patients with significant stenosis in multiple arteries were also significantly higher in the AS group (8.2% vs 1.8%, $p=0.037$). According to the study results, patients with AS are at a higher risk for MAS. Chronic mesenteric ischemia should be kept in mind in patients with AS who have symptoms such as non-specific abdominal pain and weight loss.

INTRODUCTION

Mesenteric ischemia is a rare but serious condition with high mortality rate.¹ The most common mechanism in acute cases is acute mesenteric ischemia associated with embolism. In chronic cases, mesenteric artery stenosis (MAS) associated with atherosclerosis is more common.² MAS is frequently asymptomatic due to extensive collateral network within the mesenteric vasculature. Despite that, patients with multiple stenoses and an underlying circulatory disorder such as heart failure may have lower symptom thresholds.³ Aortic stenosis (AS) is the most common type of heart valve disease in advanced age, also morbidity and mortality are quite high in symptomatic cases.⁴

The most frequent cause in adults is senile calcification. AS has pathophysiological mechanisms similar to those of atherosclerosis which

Significance of this study

What is already known about this subject?

- ▶ Mesenteric ischemia is a rare but serious condition with high mortality rate.¹ The most common mechanism in acute cases is acute mesenteric ischemia associated with embolism. In chronic cases, mesenteric artery stenosis (MAS) associated with atherosclerosis is more common.
- ▶ Aortic stenosis (AS) is the most common type of heart valve disease in advanced age, also morbidity and mortality are quite high in symptomatic cases. The most frequent cause in adults is senile calcification.
- ▶ Previous studies have shown that both coronary artery disease and peripheral artery disease are more frequent in the AS population.

What are the new findings?

- ▶ We noticed that abdominal pain complaints were common in patients hospitalized for severe AS in our clinic. In the light of these observations, we thought that the incidence of atherosclerotic stenosis would be higher in mesenteric arteries in patients with severe AS and these patients might be more symptomatic due to low flow.
- ▶ As far as we can scan, the relationship between severe AS and MAS has not been reported in the literature previously.
- ▶ In our study, the rate of patients with significant stenosis in at least 1 mesenteric artery, the rate of patients with significant stenosis in multiple vessels, and the degree of stenosis in the superior mesenteric artery and inferior mesenteric artery were significantly higher in the AS group.

involve endothelial damage, lipid accumulation, calcification, inflammation and similar risk factors. Studies have shown that both coronary artery disease and peripheral artery disease are more frequent in the AS population.^{5 6} We noticed that abdominal pain complaints were common in patients hospitalized for severe AS in our clinic. Moreover, we detected severe superior mesenteric artery (SMA) stenosis in 2 of these patients, and it was successfully treated



© American Federation for Medical Research 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Idil Soylu A, Avcioglu U, Uzunkaya F, et al. *J Investig Med* 2021;**69**:719–723.

Significance of this study

How might these results change the focus of research or clinical practice?

► In this report, the increased calcification and stenosis rate in the mesenteric arteries of an elderly and comorbidized patient group with severe AS planned for transcatheter aortic valve implantation is presented. In patients with AS, which causes a decrease in cardiac output, concomitant MAS may be the possible cause of abdominal pain, weight loss or some non-specific symptoms encountered in these patients.

percutaneously. In the light of these observations, we thought that the incidence of atherosclerotic stenosis would be higher in mesenteric arteries in patients with severe AS and these patients might be more symptomatic due to low flow. In the present study, routine CT angiography (CTA) of patients planned for transcatheter aortic valve implantation (TAVI) due to severe AS was scanned and MAS was evaluated.

MATERIALS AND METHODS

Study design and patient population

Our study was planned as a single-center, retrospective, randomized and sequential patient recruitment basis. Between 2017 and 2020, patients with aortic valve stenosis (AS group) who underwent routine CTA due to severe AS before TAVI procedure and the patients who had CTA for other indications (control group) were scanned.

Patients under 65 years of age, and those with bicuspid aortic valve, rheumatic aortic valve, hypertrophic obstructive cardiomyopathy, supraventricular AS, acute mesenteric ischemia, acute abdomen, aneurysm or dissection in the abdominal aorta, aneurysm or dissection or variation in celiac trunk (CT) or SMA, patients with no echocardiography within 1 year and patients with insufficient quality of examination were excluded from the study.

Demographic data such as gender, age, creatinine level, diabetes mellitus and dyslipidemia were recorded for each patient. Antihyperlipidemic drug use or disorder in one of the lipid values was accepted as dyslipidemia.

Image acquisition

Multidetector CTA was performed on a 64-row multidetector CT scanner (LightSpeed 64 General Electric Discovery CT750HD 2015, Milwaukee, Wisconsin, USA) with 120 kV, 300 mA, 0.98 pitch value and table speed of 64 mm per rotation. After Optiray (Dublin, Ireland), which is a 100 mL non-ionic contrast agent, was given at 3.5 mL/h, arterial phase images with a cross section thickness of 0.625 mm were obtained from the subdiaphragmatic area to the superficial femoral arteries in the 25th second.

CTA data were analyzed independently by 2 radiologists on coronal, sagittal and axial planes using OSIRIX LITE V3.8.1 (Geneva, Switzerland) program at the workstation.

CT analysis

CT, SMA and inferior mesenteric artery (IMA) stenosis rate and degree of calcification were determined in all CTA

Table 1 The classification of arterial calcification

Degree	Distribution and size of calcification
Class 0	Absent
Class 1	Single focus <5 mm and <5 foci calcification
Class 2	Single focus 5–10 mm or single focus <5 mm and ≥6 foci calcification
Class 3	Single focus >10 mm or 5–10 mm multifocal calcification

series. Arterial diameters were measured in the normal (a) and narrowest (b) lumens. The percentage of stenosis was calculated using the North American Symptomatic Carotid Endarterectomy Trial formula ($s=(a-b)/a \times 100$). At least 50% stenosis was considered as significant stenosis.

All CTAs were evaluated for calcification by the same radiologists. For this purpose, a calcification grading specific to our study was developed based on the scoring system used by van Rossum *et al*⁷ (table 1).

Statistical analysis

Statistical analysis was carried out using SPSS for Windows V.15.0. Descriptive statistics were given as mean, SD, frequency, and percentage. The Kolmogorov-Smirnov test was used to evaluate whether continuous variables were normally distributed. An independent samples t-test was used to compare mean values between the 2 groups, and χ^2 test was used for comparison of categorical data. Correlation between any 2 data was tested with the Spearman correlation analysis. Before this analysis, logarithmic transformation was applied on variables without normal distribution. Values of $p < 0.05$ were considered statistically significant.

RESULTS

A total of 246 patients (104 in the AS group and 142 in the control group) were enrolled in the study according to inclusion criteria. After exclusion criteria were applied, 184 patients were included for further analysis (73 in the AS group and 111 in the control group).

The demographic characteristics of the AS group and the control group were similar in terms of age (80.4 ± 6.7 vs 79.2 ± 5.8 , $p=0.194$), gender (54.8% vs 55% female, $p=0.983$), diabetes mellitus (21.9% vs 15.3%, $p=0.253$), dyslipidemia ratios (38% vs 42.9%, $p=0.523$) and creatinine levels (1.35 ± 1.08 mg/dL vs 1.19 ± 0.75 mg/dL, $p=0.252$).

The comparison of CT, SMA and IMA calcification scores showed that the CT calcification score was statistically similar in the AS and control groups ($p=0.148$). In contrast, class 3 calcification was never detected in the control group, while it was present in 3 patients of the AS group ($p=0.061$). SMA and IMA calcification scores were significantly higher in AS group ($p=0.035$ for SMA and $p=0.020$ for IMA). In a more detailed analysis of calcification scores, there was a more significant difference in class 3 calcification ($p=0.061$ for CT, $p=0.007$ for SMA, $p=0.031$ for IMA) (table 2).

The prevalence of significant stenosis ($\geq 50\%$) in at least 1 mesenteric artery and in 2 or 3 mesenteric arteries (multiple) among the study population was 31.5% ($n=58$) and 4.3% ($n=8$), respectively. The rate of patients with MAS in at least 1 artery (45.2% vs 22.5%, $p=0.001$) and in

Table 2 Distribution of the calcification in the mesenteric arteries

	Aortic stenosis n=73	Control n=111	P value
CT calcification score			0.148
0	28 (38.4)	51 (45.9)	0.194
1	37 (50.7)	54 (48.6)	0.452
2	5 (6.8)	6 (5.4)	0.458
3	3 (4.1)	0	0.061
SMA calcification score			0.035
0	21 (28.8)	43 (38.7)	0.165
1	35 (47.9)	56 (50.5)	0.739
2	9 (12.3)	10 (9)	0.469
3	8 (11)	2 (1.8)	0.007
IMA calcification score			0.020
0	54 (74)	97 (87.4)	0.020
1	14 (19.2)	14 (12.6)	0.225
2	2 (2.7)	0	0.080
3	3 (4.1)	0	0.031

CT, celiac trunk; IMA, inferior mesenteric artery; SMA, superior mesenteric artery.

multiple arteries (8.2% vs 1.8%, $p=0.037$) was also significantly higher in the AS group (figure 1). In addition, the rate of patients with significant MAS ($\geq 50\%$) in the AS group was higher for all 3 arteries than the control group ($p=0.05$ for CT, $p=0.025$ for SMA and $p=0.032$ for IMA). The comparison of CT, SMA and IMA stenosis degrees among the groups revealed that CT stenosis degree was similar between the groups ($p=0.322$); SMA ($p=0.025$) and IMA ($p=0.038$) stenosis degrees were higher in the AS group (table 3) (figure 2). A significant positive correlation was found between the score of calcification in each artery localization and the degree of stenosis in that artery ($r=0.430$ for CT, $p<0.001$; $r=0.534$ for SMA, $p<0.001$; and $r=0.486$ for IMA, $p<0.001$).

DISCUSSION

The most common cause of AS in adults is senile calcification. It affects 2%–3% of those over 65 years of age and up to 8% of those over 85 years. Thus, its incidence increases exponentially with age.^{8,9} The average age of patients with AS in our study was 80.4 ± 6.7 years which was higher than the average age of AS in the society. The reason for




Figure 1 Distribution of patients with significant mesenteric artery stenosis (MAS) in the 2 groups by the number of affected arteries. AS, aortic stenosis.

Table 3 Properties of the mesenteric arterial stenosis in patients

	Aortic stenosis n=73	Control n=111	P value
Critical stenosis (%)	33 (45.2)	25 (22.5)	0.001
Multivessel critical stenosis (%)	6 (8.2)	2 (1.8)	0.037
CT stenosis degree	18.2±21.2	15.4±17.6	0.322
Critical stenosis in CT (%)	11 (15.1)	7 (6.3)	0.050
SMA stenosis degree	20.3±19.2	14.9±13.4	0.025
Critical stenosis in SMA (%)	10 (13.7)	5 (4.5)	0.026
IMA stenosis degree	22.3±33.4	13.1±26.0	0.038
Critical stenosis in IMA (%)	19 (26)	15 (13.5)	0.032
Total occlusion (%)	3 (4.1)	2 (1.8)	0.346

CT, celiac trunk; IMA, inferior mesenteric artery; SMA, superior mesenteric artery.

this was that the patients included in our study consisted of patients who had high risk of surgical aortic valve replacement planned for TAVI. The control group was also scanned within a similar age range. Risk factors such as age, diabetes mellitus, hypertension and dyslipidemia that would be important for the development of atherosclerosis were found similar in both groups.

Although mesenteric ischemia constitutes only 0.1% of the admissions to the hospital, it has high mortality rate (21%–94%).^{1,10} In autopsy series, the reported range of mesenteric disease rates is from 30% to 80%.^{11,12} Atherosclerotic stenosis of the mesenteric arteries can lead to both chronic mesenteric ischemia and acute ischemia complicated by thrombosis. In 2 studies using duplex ultrasonography, significant MAS was detected in 17%–18% of patients over 65 years of age.^{13,14} In our study, this rate was found to be 31.5%. In another analysis made in patients whom renal MR angiography was performed, the rate of patients with incidental MAS was reported to be 33%.¹⁵ This heterogeneity between studies depends on both the different patient populations and the methods used in MAS detection. Since a flow rate-based calculation is used for the measurements made with duplex ultrasonography, there may be false negativity of some anatomically important stenoses. Non-critical stenoses that do not affect the flow can be visualized with MR and CT, also additional parameters such as calcification and plaque involvement properties can be evaluated. Schaefer *et al*¹⁶ compared digital subtraction angiography, duplex ultrasonography, CT and MR angiography methods




Figure 2 Distribution of patients with significant mesenteric artery stenosis (MAS) in the 2 groups according to artery localization. AS, aortic stenosis; IMA, inferior mesenteric artery; SMA, superior mesenteric artery.

in the evaluation of MAS. The researchers emphasized that the method which had the highest correlation for image quality and stenosis grading was CTA.

Plaque calcifications reflect the active phase of vascular atherosclerosis associated with inflammation. Inflammatory cytokines, such as tumor necrosis factor- α , activate osteogenic differentiation and mineralization of the extracellular matrix.¹⁷ Especially, the relationship between intimal calcification and atherosclerosis has been demonstrated in many studies.^{18–20} Moreover, calcified plaques are found to be an independent predictor of combined vascular events.^{20–21} In our study, there was a significant positive correlation between calcification score and MAS degree in all 3 arteries. In addition, calcification scores in the AS group were significantly higher in SMA and IMA localization compared with the control group. In the subgroup analysis, it was observed that the degree of calcification with the most significant difference was class 3 (large or multifocal). This suggested that multifocal and large calcifications were more common in the AS group. Depending on many factors such as size, shape, distribution in the plaque and the degree of differentiation, the effect of calcification on plaque stability may vary.¹⁸ It has been reported that multifocal calcifications increase the risk of intraplaque bleeding, provoke inflammation and trigger plaque rupture.^{19–22–23} This may be due to increased surface area and biomechanical stress in the plaque because of increased number of calcifications.

As far as we can scan, the relationship between severe AS and MAS has not been reported in the literature previously. The observation that underlies the hypothesis of this study is frequent abdominal pain symptoms that we noticed in the patients whom we admitted to our clinic to perform TAVI. Moreover, severe SMA stenosis was detected in 2 of these patients and clinical improvement was achieved after treated percutaneously. It is a well-known fact that AS and atherosclerosis have similar physiopathology because they have similar risk factors. Also, histopathological changes, such as endothelial damage, inflammation and calcification, are similar. In many studies, coronary artery disease,^{6–24} peripheral artery disease^{25–26} and carotid artery stenosis²⁷ have been found to be more frequent in patients with AS. In our study, the rate of patients with significant stenosis in at least 1 mesenteric artery or in multiple vessels, and the degree of stenosis in the SMA and IMA arteries were significantly higher in the AS group. The increased atherosclerotic risk factors of patients with AS are the probable cause of this result. Studies in the literature show that mesenteric ischemia usually occurs in patients with multiple vascular disease and severe stenosis.^{1–28} The reason for this is the intense collateral circulation in the mesenteric arterial bed.³ In contrast, circulatory disorders such as hypovolemic shock, heart failure, and myocardial infarction are known to facilitate mesenteric ischemia.² Although not previously reported, severe AS which reduces cardiac output is a good example of promoting ischemia in an already diseased mesenteric arterial bed. In addition, impaired left ventricular systolic function may increase this risk in a significant number of these patients.

Study limitations

Our study is a retrospective study and clinical symptoms of the patients that may be associated with mesenteric

ischemia have not been questioned. Therefore, the clinical reflection of the higher MAS rate in the AS group could not be analyzed in this study. In addition, although some atherosclerotic risk factors such as diabetes mellitus, hypertension, and dyslipidemia were tested, some risk factors such as smoking and heredity could not be found in most of the control group patients during file scanning. Therefore, it is not known how these factors may affect the outcome of the study. Finally, the lack of healthy individuals in the control group may have influenced the results of the study. In a study where the control group consists of healthy elderly individuals, we estimate that the results will be more significant.

CONCLUSION

In this report, the increased calcification and stenosis rate in the mesenteric arteries of an elderly and comorbidized patient group with severe AS planned for TAVI is presented. In patients with AS, which causes a decrease in cardiac output, concomitant MAS may be the possible cause of abdominal pain, weight loss or some non-specific symptoms encountered in these patients. Future prospective studies are necessary to clarify the uncertain points on this issue.

Contributors AIS: constructing the hypothesis, writing the article, saving data, image analysis. UA: establishment of hypothesis, editing of the article. FU: image analysis. KS: establishing the hypothesis, editing the article, checking the stages of the study.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Ethics approval This study was approved by the institutional ethics committee. The requirement of informed consent was waived, as this was a retrospective study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement

Data may be obtained from a third party and are not publicly available. Image data are in the hospital database. The study data can be accessed as statistical code.

ORCID iDs

Aysegül İdil Soylu <http://orcid.org/0000-0002-1390-1030>

Korhan Soylu <http://orcid.org/0000-0001-7908-2763>

REFERENCES

- 1 Thomas JH, Blake K, Pierce GE, *et al.* The clinical course of asymptomatic mesenteric arterial stenosis. *J Vasc Surg* 1998;27:840–4.
- 2 Gnanapandithan K, Feuerstadt P. Review article: mesenteric ischemia. *Curr Gastroenterol Rep* 2020;22:17.
- 3 Al-Diery H, Phillips A, Evennett N, *et al.* The pathogenesis of Nonocclusive mesenteric ischemia: implications for research and clinical practice. *J Intensive Care Med* 2019;34:771–81.
- 4 Freeman RV, Otto CM. Spectrum of calcific aortic valve disease: pathogenesis, disease progression, and treatment strategies. *Circulation* 2005;111:3316–26.
- 5 Kurra V, Schoenhagen P, Roselli EE, *et al.* Prevalence of significant peripheral artery disease in patients evaluated for percutaneous aortic valve insertion: Preprocedural assessment with multidetector computed tomography. *J Thorac Cardiovasc Surg* 2009;137:1258–64.
- 6 D'Ascenzo F, Conrotto F, Giordana F, *et al.* Mid-Term prognostic value of coronary artery disease in patients undergoing transcatheter aortic valve implantation: a meta-analysis of adjusted observational results. *Int J Cardiol* 2013;168:2528–32.
- 7 van Rossum PSN, Haverkamp L, Verkooijen HM, *et al.* Calcification of arteries supplying the gastric tube: a new risk factor for anastomotic leakage after esophageal surgery. *Radiology* 2015;274:124–32.

- 8 Stewart BF, Siscovick D, Lind BK, *et al.* Clinical factors associated with calcific aortic valve disease. cardiovascular health study. *J Am Coll Cardiol* 1997;29:630–4.
- 9 Lindroos M, Kupari M, Heikkilä J, *et al.* Prevalence of aortic valve abnormalities in the elderly: an echocardiographic study of a random population sample. *J Am Coll Cardiol* 1993;21:1220–5.
- 10 Roussel A, Castier Y, Nuzzo A, *et al.* Revascularization of acute mesenteric ischemia after creation of a dedicated multidisciplinary center. *J Vasc Surg* 2015;62:1251–6.
- 11 Reiner L, Rodriguez FL, Jimenez F, *et al.* Injection studies on mesenteric arterial circulation. III. Occlusions without intestinal infarction. *Arch Pathol* 1962;73:461–72.
- 12 Koikkalainen K, Laustela E, Tala P. Pathologic study of arteriosclerosis in the coeliac and mesenteric circulation. *Ann Chir Gynaecol Fenn* 1968;57:234–8.
- 13 Wilson DB, Mostafavi K, Craven TE, *et al.* Clinical course of mesenteric artery stenosis in elderly Americans. *Arch Intern Med* 2006;166:2095–100.
- 14 Roobottom CA, Dubbins PA. Significant disease of the celiac and superior mesenteric arteries in asymptomatic patients: predictive value of Doppler sonography. *AJR Am J Roentgenol* 1993;161:985–8.
- 15 Glockner JF. Incidental findings on renal Mr angiography. *AJR Am J Roentgenol* 2007;189:693–700.
- 16 Schaefer PJ, Pfarr J, Trentmann J, *et al.* Comparison of noninvasive imaging modalities for stenosis grading in mesenteric arteries. *Rofo* 2013;185:628–34.
- 17 Tintut Y, Patel J, Parhami F, *et al.* Tumor necrosis factor- α promotes in vitro calcification of vascular cells via the cAMP pathway. *Circulation* 2000;102:2636–42.
- 18 Shi X, Gao J, Lv Q, *et al.* Calcification in atherosclerotic plaque vulnerability: friend or foe? *Front Physiol* 2020;11:56.
- 19 Lin R, Chen S, Liu G, *et al.* Association between carotid atherosclerotic plaque calcification and intraplaque hemorrhage: a magnetic resonance imaging study. *Arterioscler Thromb Vasc Biol* 2017;37:1228–33.
- 20 Prabhakaran S, Singh R, Zhou X, *et al.* Presence of calcified carotid plaque predicts vascular events: the Northern Manhattan study. *Atherosclerosis* 2007;195:e197–201.
- 21 van den Bouwhuisen QJA, Bos D, Ikram MA, *et al.* Coexistence of calcification, intraplaque hemorrhage and lipid core within the asymptomatic atherosclerotic carotid plaque: the Rotterdam study. *Cerebrovasc Dis* 2015;39:319–24.
- 22 Nadra I, Mason JC, Philippidis P, *et al.* Proinflammatory activation of macrophages by basic calcium phosphate crystals via protein kinase C and MAP kinase pathways: a vicious cycle of inflammation and arterial calcification? *Circ Res* 2005;96:1248–56.
- 23 Mizukoshi M, Kubo T, Takarada S, *et al.* Coronary superficial and spotty calcium deposits in culprit coronary lesions of acute coronary syndrome as determined by optical coherence tomography. *Am J Cardiol* 2013;112:34–40.
- 24 Kvidal P, Bergström R, Hörte LG, *et al.* Observed and relative survival after aortic valve replacement. *J Am Coll Cardiol* 2000;35:747–56.
- 25 Sinning J-M, Horack M, Grube E, *et al.* The impact of peripheral arterial disease on early outcome after transcatheter aortic valve implantation: results from the German transcatheter aortic valve interventions registry. *Am Heart J* 2012;164:102–10.
- 26 Thomas M, Schymik G, Walther T, *et al.* Thirty-day results of the SAPIEN aortic bioprosthesis European outcome (source) registry: a European registry of transcatheter aortic valve implantation using the Edwards SAPIEN valve. *Circulation* 2010;122:62–9.
- 27 Ben-Shoshan J, Zahler D, Steinvil A, *et al.* Extracranial carotid artery stenosis and outcomes of patients undergoing transcatheter aortic valve replacement. *Int J Cardiol* 2017;227:278–83.
- 28 Bobadilla JL. Mesenteric ischemia. *Surg Clin North Am* 2013;93:925–40.