

Supplemental Table 2. Forward and reverse primers used to measure expression of adenosine receptors and cholesterol transport proteins.

Gene	Sequence
A2AR	Forward: 5'-ACCTGCAGAACGTCACCAAC-3' Reverse: 5'-TCTGCTTCAGCTGTCGTCGC-3'
A2BR	Forward: 5'-CAGACGCCACCAACTACTT-3' Reverse: 5'-GCCACCAGGAAAATCTTAATG-3'
A3R	Forward: 5'-ACTTAGCCGTCTTGA ACTCC-3' Reverse: 5'-ACCACTCACAGAAGAATATG-3'
ABCA1	Forward: 5'-GAAGTACATCAGAACATGGGC-3' Reverse: 5'-GATCAAAGCCATGGCTGTAG-3'
ABCG1	Forward: 5'-CAGGAAGATTAGACTGTGG-3' Reverse: 5'-GAAAGGGGAATGGAGAGAAG-3'
27-hydroxylase	Forward: 5'-AAGCGATACCTGGATGGTTG-3' Reverse: 5'-TGTTGGATGTCGTGTCCACT-3'
LXR- α	Forward: 5'-GGGGCCAGCCCCAAAATGCTG-3' Reverse: 5'-GCATCCGTGGGAACATCAGTCG-3'